

John Skoch Co. 1/4 Hrh3 Cas9-CKO Strategy Ronald Colons

Conplainax Ch JiaYu

Project Overview



Project Name Hrh3


Project type Cas9-CKO

Strain background C57BL/6J

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Hrh3* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Hrh3* gene has 8 transcripts. According to the structure of *Hrh3* gene, exon2 of *Hrh3-201*(ENSMUST0000056480.9) transcript is recommended as the knockout region. The region contains 167bp coding sequence.

 Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Hrh3* gene. The brief process is as follows:sgRNA was transcribed in vitro, donor vector was constructed.Cas9, sgRNA and Donor were microinjected into the fertilized eggs of C57BL/6J mice.Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6J mice.
- > The flox mice was knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- ➤ According to the existing MGI data, Homozygotes for a targeted null mutation exhibit reduced locomotor activity and body temperature, and attenuated behavioral responses to the drugs thioperamide, methamphetamine, and scopolamine.
- > The *Hrh3* gene is located on the Chr2. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Hrh3 histamine receptor H3 [Mus musculus (house mouse)]

Gene ID: 99296, updated on 12-Mar-2019

Summary

^ ?

Official Symbol Hrh3 provided by MGI

Official Full Name histamine receptor H3 provided by MGI

Primary source MGI:MGI:2139279

See related Ensembl:ENSMUSG00000039059

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as AW049250, Eae8, H3R, HH3R

Expression Biased expression in frontal lobe adult (RPKM 17.2), cortex adult (RPKM 16.0) and 8 other tissuesSee more

Orthologs <u>human</u> <u>all</u>


Transcript information (Ensembl)



The gene has 8 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Hrh3-201	ENSMUST00000056480.9	2706	<u>445aa</u>	Protein coding	CCDS17169	P58406 Q540P3	TSL:1 GENCODE basic APPRIS P2
Hrh3-202	ENSMUST00000163215.7	1401	<u>466aa</u>	Protein coding	-	E9Q540	TSL:5 GENCODE basic
Hrh3-205	ENSMUST00000165762.7	1368	<u>455aa</u>	Protein coding	-	E9Q292	TSL:5 GENCODE basic APPRIS ALT2
Hrh3-203	ENSMUST00000164442.1	1242	<u>413aa</u>	Protein coding	-	E9Q5S3	TSL:5 GENCODE basic APPRIS ALT2
Hrh3-204	ENSMUST00000165248.7	1224	<u>407aa</u>	Protein coding	-	<u>E9Q7T5</u>	TSL:5 GENCODE basic APPRIS ALT2
Hrh3-208	ENSMUST00000171736.7	906	<u>301aa</u>	Protein coding	-	E9Q522	TSL:5 GENCODE basic
Hrh3-207	ENSMUST00000166724.1	1574	<u>94aa</u>	Nonsense mediated decay	-	E9PZM9	TSL:1
Hrh3-206	ENSMUST00000166392.1	1040	No protein	Retained intron	-	-	TSL:3
	100						

The strategy is based on the design of *Hrh3-201* transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygotes for a targeted null mutation exhibit reduced locomotor activity and body temperature, and attenuated behavioral responses to the drugs thioperamide, methamphetamine, and scopolamine.



If you have any questions, you are welcome to inquire.

Tel: 025-5864 1534





