

Il16 Cas9-CKO Strategy

Designer: Baocheng Zhuang

Reviewer: Yang Zeng

Design Date: 2018-6-26

Project Overview



Project Name Il16


Project type Cas9-CKO

Strain background C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Il16* gene. The schematic diagram is as follows:



Technical routes



- The *Il16* gene has 6 transcripts. According to the structure of *Il16* gene, exon3 of *Il16-201*(ENSMUST00000001792.11) transcript is recommended as the knockout region. The region contains 109bp coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Il16* gene. The brief process is as follows:sgRNA was transcribed in vitro, donor vector was constructed.Cas9, sgRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- > The flox mice was knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- > According to the existing MGI data,mice homozygous for a knock-out allele display a transient but consistent increase of thymidine incorporation in anti-CD3-stimulated CD4+ T cells, but fail to show a hyperproliferative T cell phenotype using BrdU labeling.
- > Transcript *Il16*-202/203/204/206 may not be affected.
- > The *Il16* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



II16 interleukin 16 [Mus musculus (house mouse)]

Gene ID: 16170, updated on 13-Mar-2020

Summary

☆ ?

Official Symbol II16 provided by MGI

Official Full Name interleukin 16 provided by MGI

Primary source MGI:MGI:1270855

See related Ensembl: ENSMUSG00000001741

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as mKIAA4048

Expression Biased expression in thymus adult (RPKM 26.8), spleen adult (RPKM 17.2) and 11 other tissuesSee more

Orthologs <u>human all</u>


Transcript information (Ensembl)



The gene has 6 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
II16-201	ENSMUST00000001792.11	5260	1322aa	Protein coding	CCDS21412	<u>O54824</u>	TSL:1 GENCODE basic APPRIS is a system to annotate alternatively spliced transcripts based on a range of computational methods to identify the most functionally important transcript(s) of a gene. APPRIS P1
II16-203	ENSMUST00000145610.7	3183	624aa	Protein coding	-	<u>O54824</u>	TSL:1 GENCODE basic
II16-202	ENSMUST00000131916.1	532	29aa	Protein coding	-	D3Z3C7	CDS 3' incomplete TSL:2
II16-206	ENSMUST00000156553.7	348	<u>29aa</u>	Protein coding	9	D3Z3C7	CDS 3' incomplete TSL:2
II16-205	ENSMUST00000153560.1	2625	238aa	Nonsense mediated decay		D6RI15	TSL:1
1116-204	ENSMUST00000151047.1	2605	No protein	Retained intron	-	745	TSL:1

The strategy is based on the design of *Il16-201* transcript, the transcription is shown below:




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data,mice homozygous for a knock-out allele display a transient but consistent increase of thymidine incorporation in anti-CD3-stimulated CD4+ T cells, but fail to show a hyperproliferative T cell phenotype using BrdU labeling.



If you have any questions, you are welcome to inquire.

Tel: 025-5864 1534





