

Fcgrt Cas9-KO Strategy

Designer:

Reviewer:

Design Date:


Xiaojing Li

2020-4-8

Jing Jin

Project Overview






Knockout strategy



400-9660890

This model will use CRISPR/Cas9 technology to edit the *Fcgrt* gene. The schematic diagram is as follows:



江苏集萃药康生物科技股份有限公司

GemPharmatech Co., Ltd.



> The *Fcgrt* gene has 1 transcripts. According to the structure of *Fcgrt* gene, exon2-exon6 of MGP_BALBcJ_T0083760.1 transcript is recommended as the knockout region. The region contains start codon ATG.Knock out the region will result in disruption of protein function.

In this project we use CRISPR/Cas9 technology to modify *Fcgrt* gene. The brief process is as follows: CRISPR/Cas9 system transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with BALB/C mice.

- According to the existing MGI data, Homozygous mutation of this gene results in defective perinatal transport of maternal IgG, increased clearance of IgG, and diminished IgG antibody response after immunization.
- The KO region contains functional region of the Fcgrt gene.Knockout the region may affect the function of Rcn3 gene.
 The *Fcgrt* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Notice

Gene information (NCBI)



☆ ?

Fcgrt Fc receptor, IgG, alpha chain transporter [Mus musculus (house mouse)]

Gene ID: 14132, updated on 13-Mar-2020

- Summary

Official SymbolFcgrt provided by MGIOfficial Full NameFc receptor, IgG, alpha chain transporter provided by MGIPrimary sourceMGI:MGI:103017See relatedEnsembl:ENSMUSG0000003420Gene typeprotein codingRefSeq statusVALIDATEDOrganismMus musculusLineageEukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;
Muroidea; Muridae; Murinae; Mus; MusAlso known asFcRnExpressionBroad expression in placenta adult (RPKM 121.9), mammary gland adult (RPKM 87.9) and 23 other tissuesSee more
human all

江苏集萃药康生物科技股份有限公司

GemPharmatech Co., Ltd.


Transcript information (Ensembl)



The gene has 1 transcripts, all transcripts are shown below:

Name 🍦	Transcript ID 🖕	bp 🌲	Protein 🖕	Biotype 🍦	CCDS 🖕	UniProt 🖕	Flags
1	MGP_BALBcJ_T0083760.1	1738	<u>365aa</u>	Protein coding	<u>CCDS52243</u> &	<u>Q61559</u> & <u>Q6PKB0</u> &	2

The strategy is based on the design of MGP_BALBcJ_T0083760.1 transcript, The transcription is shown below:




江苏集萃药康生物科技股份有限公司

GemPharmatech Co., Ltd.

Genomic location distribution





江苏集萃药康生物科技股份有限公司

GemPharmatech Co., Ltd.

Protein domain




	· 3	(0	1975		VO2	Ŷ	2		-	(AX	
7.3.	MGP_BALBcJ_P0083 Transmembrane heli Low complexity (Seg) Cleavage site (Sign Superfamily	MHC	lasses I/II-like an	itigen recognition p	rotein		Immunoglol	pulin-like domain su	uperfamily			5 Co.: <
	SMART						Imm	unoglobulin C1-set	_			
	Pfam.	MHC	class I-like antigen	recognition-like				noglobulin C1-set	1			
	PROSITE profiles							bulin-like domain	_			
	PROSITE patterns						Immunogio	pulin-like domain	Immunoglob	ulin/major histocor	npatibility cor	0
	PANTHER	THR16675:SF3									_	- 2
	Gene3D	PTHR16675 MHC class I-like antigen recognition-like superfamily					Immunoglob	Immunoglobulin-like fold				
	Scale bar 0		40	80	120	160	200	240	280	320	365	1
	Ad Marker		narm.				Deharman Pharman			narma.	× × ×	
	~?~			0	,	\sim		So.				

江苏集萃药康生物科技股份有限公司

GemPharmatech Co., Ltd.

Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous mutation of this gene results in defective perinatal transport of maternal IgG, increased clearance of IgG, and diminished IgG antibody response after immunization.



If you have any questions, you are welcome to inquire. Tel: 400-9660890



