

# *Hpgd* Cas9-KO Strategy

Designer: Daohua Xu

# Project Overview

---

**Project Name**

*Hpgd*

---

**Project type**

**Cas9-KO**

---


**Strain background**

**C57BL/6JGpt**

---

# Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Hpgd* gene. The schematic diagram is as follows:



- The *Hpgd* gene has 1 transcript. According to the structure of *Hpgd* gene, exon3 of *Hpgd-201*(ENSMUST00000034026.10) transcript is recommended as the knockout region. The region contains 107bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Hpgd* gene. The brief process is as follows: CRISPR/Cas9 system were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.

- According to the existing MGI data, homozygous mutation of this gene results failure of the ductus arteriosus to close and perinatal lethality. Mutant animals die within 12-48 hours after birth due to congestive heart failure. Mice homozygous for a hypomorphic allele exhibit preterm labor.
- The *Hpgd* gene is located on the Chr8. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.



# Gene information (NCBI)

## Hpgd hydroxyprostaglandin dehydrogenase 15 (NAD) [Mus musculus (house mouse)]

Gene ID: 15446, updated on 13-Mar-2020

### Summary



**Official Symbol** Hpgd provided by [MGI](#)

**Official Full Name** hydroxyprostaglandin dehydrogenase 15 (NAD) provided by [MGI](#)

**Primary source** [MGI:MGI:108085](#)

**See related** [Ensembl:ENSMUSG00000031613](#)

**Gene type** protein coding

**RefSeq status** VALIDATED

**Organism** [Mus musculus](#)

**Lineage** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

**Also known as** 15-PGDH, AV026552

**Expression** Broad expression in lung adult (RPKM 76.0), bladder adult (RPKM 50.8) and 17 other tissues [See more](#)


**Orthologs** [human](#) [all](#)

# Transcript information (Ensembl)


The gene has 1 transcript,and the transcript is shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Hpgd-201	<a href="#">ENSMUST00000034026.9</a>	1683	<a href="#">269aa</a>	Protein coding	<a href="#">CCDS40341</a>	<a href="#">Q8VCC1</a>	TSL:1 GENCODE basic APPRIS is a system to annotate alternatively spliced transcripts based on a range of computational methods to identify the most functionally important transcript(s) of a gene. APPRIS P1

The strategy is based on the design of *Hpgd-201* transcript,the transcription is shown below:




# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI )



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, homozygous mutation of this gene results failure of the ductus arteriosus to close and perinatal lethality. Mutant animals die within 12-48 hours after birth due to congestive heart failure. Mice homozygous for a hypomorphic allele exhibit preterm labor.

If you have any questions, you are welcome to inquire.  
Tel: 400-9660890

