

Prdm1 Cas9-CKO Strategy

Designer: Daohua Xu

Project Overview




| Project Name | Prdm1 | | | |
|-------------------|-------------|--|--|--|
| Project type | Cas9-CKO | | | |
| Strain background | C57BL/6JGpt | | | |

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Prdm1* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Prdm1* gene has 4 transcripts. According to the structure of *Prdm1* gene, exon3-exon4 of *Prdm1*202(ENSMUST00000105490.3) transcript is recommended as the knockout region. The region contains 373bp coding sequence.

 Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Prdm1* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- > The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- ➤ According to the existing MGI data,homozygous null mice display embryonic lethality and impaired primordial germ cell development, while heterozygotes display a decreased numbers of primordial germ cells but normal migration.

 Conditional mutants display impaired plasma cell and pre-plasmamemory B cell development.
- > The *Prdm1* gene is located on the Chr10. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- ➤ This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Prdm1 PR domain containing 1, with ZNF domain [Mus musculus (house mouse)]

Gene ID: 12142, updated on 21-Feb-2021

Summary

☆ ?

Official Symbol Prdm1 provided by MGI

Official Full Name PR domain containing 1, with ZNF domain provided by MGI

Primary source MGI:MGI:99655

See related Ensembl: ENSMUSG00000038151

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as Bli, Blim, Blimp-1, Blimp1, PRDI-, PRDI-BF1, ZNFPR1A1, b2b1765C, b2b1765Clo

Expression Broad expression in placenta adult (RPKM 3.0), subcutaneous fat pad adult (RPKM 1.7) and 20 other tissuesSee more

Orthologs <u>human all</u>


Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:


| Name | Transcript ID | bp | Protein | Biotype | CCDS | UniProt | Flags |
|-----------|-----------------------|------|--------------|-----------------|-----------|---------|---------------------------------------|
| Prdm1-202 | ENSMUST00000105490.3 | 4025 | <u>823aa</u> | Protein coding | CCDS23825 | | TSL:1 , GENCODE basic , APPRIS P2 , |
| Prdm1-201 | ENSMUST00000039174.11 | 5281 | <u>856aa</u> | Protein coding | - | | TSL:2, GENCODE basic, APPRIS ALT2, |
| Prdm1-204 | ENSMUST00000218369.2 | 2517 | <u>838aa</u> | Protein coding | <u> </u> | | TSL:5 , GENCODE basic , APPRIS ALT2 , |
| Prdm1-203 | ENSMUST00000167340.2 | 830 | No protein | Retained intron | - | | TSL:1, |

The strategy is based on the design of *Prdm1-202* transcript, the transcription is shown below:




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, homozygous null mice display embryonic lethality and impaired primordial germ cell development, while heterozygotes display a decreased numbers of primordial germ cells but normal migration.

Conditional mutants display impaired plasma cell and pre-plasmamemory B cell development.



If you have any questions, you are welcome to inquire.

Tel: 400-9660890





