

# ***Fbxw8 Cas9-CKO Strategy***

**Designer: Jiayuan Yao**

**Reviewer: Lingyan Wu**

**Design Date: 2020-7-2**

# Project Overview

**Project Name**

*Fbxw8*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Fbxw8* gene. The schematic diagram is as follows:



- The *Fbxw8* gene has 2 transcripts. According to the structure of *Fbxw8* gene, exon2-exon4 of *Fbxw8*-201(ENSMUST00000049474.8) transcript is recommended as the knockout region. The region contains 359bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Fbxw8* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, mice homozygous for a null allele display partial late embryonic lethality with embryonic growth retardation and abnormal placental morphology.
- The *Fbxw8* gene is located on the Chr5. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

Fbxw8 F-box and WD-40 domain protein 8 [Mus musculus (house mouse)]

Gene ID: 231672, updated on 13-Mar-2020

## Summary



Official Symbol Fbxw8 provided by [MGI](#)

Official Full Name F-box and WD-40 domain protein 8 provided by [MGI](#)

Primary source [MGI:MGI:1923041](#)

See related [Ensembl:ENSMUSG00000032867](#)

Gene type protein coding

RefSeq status VALIDATED

Organism [Mus musculus](#)

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 4930438M06Rik, FBW6, FBW8, FBX029, Fbx29

Expression Ubiquitous expression in ovary adult (RPKM 39.6), subcutaneous fat pad adult (RPKM 36.4) and 28 other tissues [See more](#)


Orthologs [human all](#)

# Transcript information (Ensembl)


The gene has 2 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Fbxw8-201	<a href="#">ENSMUST00000049474.8</a>	5000	<a href="#">598aa</a>	Protein coding	<a href="#">CCDS19609</a>	<a href="#">Q8BIA4</a>	TSL:1 GENCODE basic APPRIS P1
Fbxw8-202	<a href="#">ENSMUST00000201545.1</a>	1937	No protein	Retained intron	-	-	TSL:1

The strategy is based on the design of *Fbxw8-201* transcript,the transcription is shown below:




# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, mice homozygous for a null allele display partial late embryonic lethality with embryonic growth retardation and abnormal placental morphology.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

