

Lhfpl4 Cas9-KO Strategy

Designer: Lingyan Wu

Reviewer: Miaomiao Cui

Design Date: 2021-3-1

Project Overview



Project Name

Lhfpl4

Project type

Cas9-KO


Strain background

C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Lhfpl4* gene. The schematic diagram is as follows:



Technical routes



- > The *Lhfpl4* gene has 2 transcripts. According to the structure of *Lhfpl4* gene, exon2 of *Lhfpl4*-201(ENSMUST00000162280.1) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Lhfpl4* gene. The brief process is as follows: CRISPR/Cas9 system were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.

Notice



- > According to the existing MGI data, homozygous knockout affects inhibitory postsynaptic currents in the hippocampus.
- > The *Lhfpl4* gene is located on the Chr6. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- > This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Lhfpl4 lipoma HMGIC fusion partner-like protein 4 [Mus musculus (house mouse)]

Gene ID: 269788, updated on 13-Mar-2020

Summary

☆ ?

Official Symbol Lhfpl4 provided by MGI

Official Full Name lipoma HMGIC fusion partner-like protein 4 provided by MGI

Primary source MGI:MGI:3057108

See related Ensembl: ENSMUSG00000042873

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 1190004M23Rik, Al604880, B230384L07, mKIAA4027

Expression Biased expression in whole brain E14.5 (RPKM 16.6), CNS E18 (RPKM 16.3) and 6 other tissuesSee more

Orthologs <u>human all</u>


Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


| Name | Transcript ID | bp | Protein | Biotype | CCDS | UniProt | Flags |
|------------|----------------------|------|--------------|----------------|-----------|------------|-------------------------------|
| Lhfpl4-201 | ENSMUST00000162280.1 | 4762 | 247aa | Protein coding | CCDS39591 | Q5U4E0 | TSL:1 GENCODE basic APPRIS P1 |
| Lhfpl4-202 | ENSMUST00000203665.1 | 1414 | <u>168aa</u> | Protein coding | 2 | A0A0N4SV06 | CDS 5' incomplete TSL:3 |

The strategy is based on the design of *Lhfpl4-201* transcript, the transcription is shown below:




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, homozygous knockout affects inhibitory postsynaptic currents in the hippocampus.



If you have any questions, you are welcome to inquire.

Tel: 400-9660890





