

# *Slc17a2* Cas9-CKO Strategy

**Designer: Lingyan Wu**

**Reviewer: Miaomiao Cui**

**Design Date: 2021-7-9**

# Project Overview

---

**Project Name**

*Slc17a2*

---

**Project type**

**Cas9-CKO**

---


**Strain background**

**C57BL/6JGpt**

---

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Slc17a2* gene. The schematic diagram is as follows:



# Technical routes

- The *Slc17a2* gene has 4 transcripts. According to the structure of *Slc17a2* gene, exon3 of *Slc17a2*-201(ENSMUST00000006786.11) transcript is recommended as the knockout region. The region contains 212bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Slc17a2* gene. The brief process is as follows: sgRNA was transcribed in vitro, donor was constructed. Cas9, sgRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice was knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.



- The *Slc17a2* gene is located on the Chr13. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

# Gene information (NCBI)

## Slc17a2 solute carrier family 17 (sodium phosphate), member 2 [Mus musculus (house mouse)]

Gene ID: 218103, updated on 17-Feb-2021

### Summary



**Official Symbol** Slc17a2 provided by [MGI](#)

**Official Full Name** solute carrier family 17 (sodium phosphate), member 2 provided by [MGI](#)

**Primary source** [MGI:MGI:2443098](#)

**See related** [Ensembl:ENSMUSG00000036110](#)

**Gene type** protein coding

**RefSeq status** VALIDATED

**Organism** [Mus musculus](#)

**Lineage** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

**Also known as** C730032N17Rik, NP, NPT3

**Expression** Biased expression in liver adult (RPKM 29.3), liver E18 (RPKM 6.3) and 1 other tissue [See more](#)


**Orthologs** [human](#) [all](#)

# Transcript information (Ensembl)

The gene has 4 transcripts,all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Slc17a2-201	<a href="#">ENSMUST00000006786.11</a>	4629	<a href="#">447aa</a>	Protein coding	<a href="#">CCDS36620</a>		TSL:1 , GENCODE basic , APPRIS P2 ,
Slc17a2-202	<a href="#">ENSMUST00000099697.3</a>	1893	<a href="#">478aa</a>	Protein coding	-		TSL:5 , GENCODE basic , APPRIS ALT2 ,
Slc17a2-203	<a href="#">ENSMUST00000139572.2</a>	3863	No protein	Retained intron	-		TSL:5 ,
Slc17a2-204	<a href="#">ENSMUST00000141717.8</a>	3269	No protein	Retained intron	-		TSL:1 ,

The strategy is based on the design of *Slc17a2-201* transcript,the transcription is shown below:






# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI )



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

If you have any questions, you are welcome to inquire.

Tel: 025-5864 1534

