

# *Oxtr* Cas9-CKO Strategy

Designer: Yun Li  
Reviewer: Longyun Hu  
Design Date: 2019-7-19

# Project Overview

**Project Name**

*Oxtr*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Oxtr* gene. The schematic diagram is as follows:



- The *Oxtr* gene has 2 transcripts. According to the structure of *Oxtr* gene, exon1 of *Oxtr-201* (ENSMUST00000053306.7) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Oxtr* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Homozygous null females fail to lactate and exhibit decreased maternal behavior. Males exhibit an increase in aggression, hypoactivity and vocalization in response to social isolation, and social amnesia.
- The *Oxtr* gene is located on the Chr6. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Oxtr oxytocin receptor [Mus musculus (house mouse)]

Gene ID: 18430, updated on 12-Mar-2019

### Summary




<b>Official Symbol</b>	Oxtr provided by <a href="#">MGI</a>
<b>Official Full Name</b>	oxytocin receptor provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:109147</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG00000049112</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	OTR
<b>Expression</b>	Biased expression in adrenal adult (RPKM 71.9), mammary gland adult (RPKM 22.4) and 3 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 2 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Oxtr-201	<a href="#">ENSMUST00000053306.7</a>	4704	<a href="#">388aa</a>	Protein coding	<a href="#">CCDS39588</a>	<a href="#">P97926</a>	TSL:1 GENCODE basic APPRIS P1
Oxtr-202	<a href="#">ENSMUST00000204027.1</a>	3721	<a href="#">75aa</a>	Protein coding	-	<a href="#">A0A0N4SVY6</a>	TSL:1 GENCODE basic

The strategy is based on the design of *Oxtr-201* transcript,The transcription is shown below



# Genomic location distribution





# Protein domain

ENSMUSP00000051...

Transmembrane heli...

MobiDB lite

Low complexity (Seg)

Conserved Domains

hmmpanther

Superfamily domains

Prints domain

Pfam domain

PROSITE profiles


PROSITE patterns

Gene3D


All sequence SNPs/i...

Variant Legend

Scale bar



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Homozygous null females fail to lactate and exhibit decreased maternal behavior.

Males exhibit an increase in aggression, hypoactivity and vocalization in response to social isolation, and social amnesia.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

