



Cdc20 Cas9-CKO Strategy

Designer:
Reviewer:
Design Date:

Huan Wang
Huan Fan
2019-11-7

Project Overview


Project Name**Cdc20**

Project type**Cas9-CKO**

Strain background**C57BL/6JGpt**

Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Cdc20* gene. The schematic diagram is as follows:



Technical routes

- The *Cdc20* gene has 3 transcripts. According to the structure of *Cdc20* gene, exon1-exon8 of *Cdc20-201* (ENSMUST00000006565.12) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cdc20* gene. The brief process is as follows:gRNA was transcribed in vitro, donor was constructed.Cas9, gRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice.Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.



集萃药康
GemPharmatech

Notice

- According to the existing MGI data, Mice homozygous for mutations in this gene display embryonic lethality with cell cycle abnormalities. Mice heterozygous for a mutation in this gene display increased tumor incidence and increased incidence of aneuploidy.
- The *Cdc20* gene is located on the Chr4. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



集萃药康
GemPharmatech

Gene information (NCBI)

Cdc20 cell division cycle 20 [Mus musculus (house mouse)]

Gene ID: 107995, updated on 5-Feb-2019

Summary



Official Symbol Cdc20 provided by [MGI](#)

Official Full Name cell division cycle 20 provided by [MGI](#)

Primary source [MGI:MGI:1859866](#)

See related [Ensembl:ENSMUSG00000006398](#)

Gene type protein coding

RefSeq status VALIDATED

Organism [Mus musculus](#)

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 2310042N09Rik, C87100, p55CDC

Expression Broad expression in liver E14.5 (RPKM 69.5), liver E14 (RPKM 66.0) and 17 other tissues [See more](#)


Orthologs [human](#) [all](#)

Transcript information (Ensembl)

The gene has 3 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Cdc20-201	ENSMUST00000006565.12	1793	499aa	Protein coding	CCDS18551	Q9JJ66	TSL:1 GENCODE basic APPRIS P1
Cdc20-203	ENSMUST00000183942.1	771	197aa	Nonsense mediated decay	-	V9GXW0	CDS 5' incomplete TSL:3
Cdc20-202	ENSMUST00000151302.1	526	No protein	Retained intron	-	-	TSL:2


The strategy is based on the design of *Cdc20-201* transcript, The transcription is shown below






集萃药康
GemPharmatech


Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous for mutations in this gene display embryonic lethality with cell cycle abnormalities. Mice heterozygous for a mutation in this gene display increased tumor incidence and increased incidence of aneuploidy.



If you have any questions, you are welcome to inquire.

Tel: 400-9660890



集萃药康生物科技
GemPharmatech Co.,Ltd

