

Cd5l Cas9-CKO Strategy

| | |
|---------------------|------------------|
| Designer: | Huan Wang |
| Reviewer: | Huan Fan |
| Design Date: | 2020-5-12 |

Project Overview

Project Name

Cd5l

Project type


Cas9-CKO

Strain background

C57BL/6JGpt

Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Cd5l* gene. The schematic diagram is as follows:



- The *Cd5l* gene has 1 transcript. According to the structure of *Cd5l* gene, exon2-exon6 of *Cd5l-201* (ENSMUST00000015998.7) transcript is recommended as the knockout region. The region contains most coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cd5l* gene. The brief process is as follows: gRNA was transcribed in vitro, donor was constructed. Cas9, gRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, homozygous null mice exhibit enhanced and prolonged bacterial-induced granuloma formation and increased apoptosis of nkt and t cells.
- The *Cd51* gene is located on the Chr3. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)

Cd5l CD5 antigen-like [Mus musculus (house mouse)]

Gene ID: 11801, updated on 13-Mar-2020

Summary



Official Symbol Cd5l provided by [MGI](#)

Official Full Name CD5 antigen-like provided by [MGI](#)

Primary source [MGI:MGI:1334419](#)

See related [Ensembl:ENSMUSG00000015854](#)

Gene type protein coding

RefSeq status VALIDATED

Organism [Mus musculus](#)

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 1/6, AAC-11, AI047839, AIM, Api6, CT2, Pdp, Sp-alpha, mAIM

Expression Biased expression in liver E18 (RPKM 55.6), liver E14 (RPKM 35.5) and 4 other tissues [See more](#)


Orthologs [human](#) [all](#)

Transcript information (Ensembl)


The gene has 1 transcript, and the transcript is shown below:

| Name | Transcript ID | bp | Protein | Biotype | CCDS | UniProt | Flags |
|----------|--------------------------------------|------|-----------------------|----------------|---------------------------|------------------------|-------------------------------|
| Cd5l-201 | ENSMUST00000015998.7 | 2036 | 352aa | Protein coding | CCDS17451 | Q9QWK4 | TSL:1 GENCODE basic APPRIS P1 |


The strategy is based on the design of *Cd5l-201* transcript, the transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, homozygous null mice exhibit enhanced and prolonged bacterial-induced granuloma formation and increased apoptosis of NKT and T cells.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

