

# *Cd33* Cas9-CKO Strategy

**Designer:**

**Huan Fan**

**Reviewer:**

**Huan Wang**

**Design Date:**

**2020-1-13**

# Project Overview

**Project Name**

*Cd33*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Cd33* gene. The schematic diagram is as follows:



# Technical routes

- The *Cd33* gene has 6 transcripts. According to the structure of *Cd33* gene, exon1-exon3 of *Cd33-203* (ENSMUST00000205503.1) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cd33* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Mice homozygous for disruptions in this gene show slight reductions in mean erythrocyte count and hematocrit and increased concentration of blood aspartate aminotransaminase. There is also a hyporesponsiveness to induced peritonitis and a weaker IL-6 response to LPS-induced systemic inflammation.
- The *Cd33* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Cd33 CD33 antigen [Mus musculus (house mouse)]

Gene ID: 12489, updated on 31-Jan-2019

### Summary




|                           |   |
|---------------------------|---|
| <b>Official Symbol</b>    | Cd33 provided by <a href="#">MGI</a>  |
| <b>Official Full Name</b> | CD33 antigen provided by <a href="#">MGI</a>  |
| <b>Primary source</b>     | <a href="#">MGI:MGI:99440</a>   |
| <b>See related</b>        | <a href="#">Ensembl:ENSMUSG000000004609</a>   |
| <b>Gene type</b>          | protein coding  |
| <b>RefSeq status</b>      | VALIDATED   |
| <b>Organism</b>           | <a href="#">Mus musculus</a>  |
| <b>Lineage</b>            | Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus |
| <b>Also known as</b>      | Siglec-3, gp67  |
| <b>Expression</b>         | Broad expression in liver E18 (RPKM 4.3), mammary gland adult (RPKM 4.2) and 27 other tissues <a href="#">See more</a>  |
| <b>Orthologs</b>          | <a href="#">human</a> <a href="#">all</a>   |

# Transcript information (Ensembl)


The gene has 6 transcripts,all transcripts are shown below:

| Name     | Transcript ID                         | bp   | Protein               | Biotype         | CCDS                      | UniProt                    | Flags                           |
|----------|---------------------------------------|------|-----------------------|-----------------|---------------------------|----------------------------|---------------------------------|
| Cd33-203 | <a href="#">ENSMUST00000205503.1</a>  | 5728 | <a href="#">334aa</a> | Protein coding  | <a href="#">CCDS21173</a> | <a href="#">Q63994</a>     | TSL:1 GENCODE basic APPRIS P3   |
| Cd33-201 | <a href="#">ENSMUST00000004728.11</a> | 1962 | <a href="#">403aa</a> | Protein coding  | <a href="#">CCDS52224</a> | <a href="#">Q63994</a>     | TSL:1 GENCODE basic APPRIS ALT2 |
| Cd33-202 | <a href="#">ENSMUST00000039861.6</a>  | 1791 | <a href="#">334aa</a> | Protein coding  | <a href="#">CCDS21173</a> | <a href="#">Q63994</a>     | TSL:1 GENCODE basic APPRIS P3   |
| Cd33-205 | <a href="#">ENSMUST00000206371.1</a>  | 933  | <a href="#">240aa</a> | Protein coding  | -                         | <a href="#">A0A0U1RNJ9</a> | CDS 5' incomplete TSL:5         |
| Cd33-204 | <a href="#">ENSMUST00000205687.1</a>  | 2376 | No protein            | Retained intron | -                         | -                          | TSL:1                           |
| Cd33-206 | <a href="#">ENSMUST00000206977.1</a>  | 402  | No protein            | lncRNA          | -                         | -                          | TSL:2                           |

The strategy is based on the design of *Cd33-203* transcript,The transcription is shown below



# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Mice homozygous for disruptions in this gene show slight reductions in mean erythrocyte count and hematocrit and increased concentration of blood aspartate aminotransaminase. There is also a hyporesponsiveness to induced peritonitis and a weaker IL-6 response to LPS-induced systemic inflammation.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

