

# Fads2 Cas9-CKO Strategy

**Designer:** Huan Wang

**Reviewer:** Huan Fan

**Design Date:** 2020-5-26

# **Project Overview**



Project Name Fads2


Project type Cas9-CKO

Strain background C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Fads2* gene. The schematic diagram is as follows:



### Technical routes



- ➤ The *Fads2* gene has 3 transcripts. According to the structure of *Fads2* gene, exon2-exon10 of *Fads2-201* (ENSMUST00000025567.8) transcript is recommended as the knockout region. The region contains 950bp coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Fads2* gene. The brief process is as follows:gRNA was transcribed in vitro, donor was constructed.Cas9, gRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice.Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

### **Notice**



- ➤ According to the existing MGI data, mice homozygous for a null allele display absence of long-chain polyunsaturated fatty acids, infertility, arrest of spermiogenesis and folliculogenesis, and impaired platelet function.
- The *Fads2* gene is located on the Chr19. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- > This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

### Gene information (NCBI)



#### Fads2 fatty acid desaturase 2 [Mus musculus (house mouse)]

Gene ID: 56473, updated on 13-Mar-2020

#### Summary

☆ ?

Official Symbol Fads2 provided by MGI

Official Full Name fatty acid desaturase 2 provided by MGI

Primary source MGI:MGI:1930079

See related Ensembl: ENSMUSG00000024665

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 2900042M13Rik, Fads2a, Fadsd2

Expression Broad expression in adrenal adult (RPKM 555.1), liver adult (RPKM 281.5) and 21 other tissuesSee more

Orthologs <u>human all</u>


# Transcript information (Ensembl)



The gene has 3 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Fads2-201	ENSMUST00000025567.8	3150	444aa	Protein coding	CCDS29571	Q9Z0R9	TSL:1 GENCODE basic APPRIS P1
Fads2-202	ENSMUST00000235937.1	355	<u>118aa</u>	Protein coding	-	A0A494BBD1	CDS 5' and 3' incomplete
Fads2-203	ENSMUST00000238023.1	879	No protein	Processed transcript	\$20	2	

The strategy is based on the design of *Fads2-201* transcript, the transcription is shown below:




### Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data,mice homozygous for a null allele display absence of long-chain polyunsaturated fatty acids, infertility, arrest of spermiogenesis and folliculogenesis, and impaired platelet function.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





