

Dhx9 Cas9-CKO Strategy

Designer:

Huan Fan

Reviewer:

Huan Wang

Design Date:

2020-2-28

Project Overview



Project Name Dhx9


Project type Cas9-CKO

Strain background C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Dhx9* gene. The schematic diagram is as follows:



Technical routes



- The *Dhx9* gene has 9 transcripts. According to the structure of *Dhx9* gene, exon3-exon4 of *Dhx9-202* (ENSMUST00000186380.6) transcript is recommended as the knockout region. The region contains 259bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Dhx9* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- ➤ According to the existing MGI data, Homozygotes die in embryonic stages with massive apoptotic cell death in embryonic ectodermal cells.
- \Rightarrow The *Dhx9* gene is located on the Chr1. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Dhx9 DEAH (Asp-Glu-Ala-His) box polypeptide 9 [Mus musculus (house mouse)]

Gene ID: 13211, updated on 31-Jan-2019

Summary

☆ ?

Official Symbol Dhx9 provided by MGI

Official Full Name DEAH (Asp-Glu-Ala-His) box polypeptide 9 provided by MGI

Primary source MGI:MGI:108177

See related Ensembl: ENSMUSG00000042699

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as Al326842, Ddx9, HEL-5, NDHII, RHA, mHEL-5

Expression Broad expression in CNS E11.5 (RPKM 71.6), CNS E14 (RPKM 55.5) and 21 other tissuesSee more

Orthologs <u>human</u> all


Transcript information (Ensembl)



The gene has 9 transcripts, all transcripts are shown below:


	No. of the contract of the con					- 1	
Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Dhx9-202	ENSMUST00000186380.6	4561	1383aa	Protein coding	CCDS78706	A0A087WPL5	TSL:2 GENCODE basic APPRIS P2
Dhx9-201	ENSMUST00000042141.11	4616	1384aa	Protein coding	6.70	E9QNN1	TSL:5 GENCODE basic APPRIS ALT2
Dhx9-204	ENSMUST00000188345.1	2346	<u>735aa</u>	Protein coding	140	Q3UR42	CDS 3' incomplete TSL:1
Dhx9-203	ENSMUST00000186966.6	2406	<u>459aa</u>	Nonsense mediated decay	VSV	A0A0R4J2C3	TSL:1
Dhx9-209	ENSMUST00000190544.6	2018	<u>47aa</u>	Nonsense mediated decay	1783	A0A087WRT3	CDS 5' incomplete TSL:1
Dhx9-206	ENSMUST00000189509.1	2658	No protein	Retained intron	6.70	-	TSL:2
Dhx9-208	ENSMUST00000190383.1	2274	No protein	Retained intron	740		TSL:NA
Dhx9-205	ENSMUST00000189425.1	1604	No protein	Retained intron	V-31	92	TSL:1
Dhx9-207	ENSMUST00000189516.1	658	No protein	Retained intron	1783	-	TSL:2
	1111	7.7	1		No. of Long.	A. Village	7 ; .

The strategy is based on the design of Dhx9-202 transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygotes die in embryonic stages with massive apoptotic cell death in embryonic ectodermal cells.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





