# *Nck2* Cas9-CKO Strategy

**Designer:** Huan Fan
**Reviewer:** Huan Wang
**Design Date:** 2020-2-28

集萃药康 GemPharmatech

# Project Overview

| | |
|---|---|
| **Project Name** | ***Nck2*** |
| **Project type** | **Cas9-CKO** |
| **Strain background** | **C57BL/6JGpt** |

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Nck2* gene. The schematic diagram is as follows:



# Technical routes

- The *Nck2* gene has 6 transcripts. According to the structure of *Nck2* gene, exon3-exon4 of *Nck2-201* (ENSMUST00000086421.8) transcript is recommended as the knockout region.The region contains start codon ATG. Knock out the region will result in disruption of protein function.

- In this project we use CRISPR/Cas9 technology to modify *Nck2* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice.Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.

- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

# Notice

- According to the existing MGI data,Mice homozygous for disruption of this gene display no abnormal phenotype.

- The *Nck2* gene is located on the Chr1. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.

- This Strategy is designed based on genetic information in existing databases.Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

# Gene information （NCBI）

集萃药康 GemPharmatech

## Nck2 non-catalytic region of tyrosine kinase adaptor protein 2 [Mus musculus (house mouse)]

Gene ID: 17974, updated on 31-Jan-2019

### Summary

| | |
|---|---|
| **Official Symbol** | Nck2 provided by MGI |
| **Official Full Name** | non-catalytic region of tyrosine kinase adaptor protein 2 provided by MGI |
| **Primary source** | MGI:MGI:1306821 |
| **See related** | Ensembl:ENSMUSG00000066877 |
| **Gene type** | protein coding |
| **RefSeq status** | VALIDATED |
| **Organism** | Mus musculus |
| **Lineage** | Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus |
| **Also known as** | 4833426I10Rik, Grb4, NCKbeta, Nck-2 |
| **Expression** | Ubiquitous expression in thymus adult (RPKM 33.3), colon adult (RPKM 26.2) and 27 other tissues See more |
| **Orthologs** | human all |

# Transcript information（Ensembl）

The gene has 6 transcripts,all transcripts are shown below:

| Name | Transcript ID | bp | Protein | Biotype | CCDS | UniProt | Flags |
|---|---|---|---|---|---|---|---|
| **Nck2-201** | ENSMUST00000086421.8 | 2772 | 380aa | Protein coding | CCDS14924 | O55033 | TSL:1 GENCODE basic APPRIS P1 |
| **Nck2-206** | ENSMUST00000202540.1 | 578 | 142aa | Protein coding | - | A0A0J9YUK4 | CDS 3' incomplete TSL:3 |
| **Nck2-202** | ENSMUST00000114744.2 | 420 | 77aa | Protein coding | - | A0A0A0MQF0 | CDS 5' incomplete TSL:5 |
| **Nck2-205** | ENSMUST00000187435.6 | 372 | 21aa | Protein coding | - | A0A087WQT7 | CDS 3' incomplete TSL:5 |
| **Nck2-204** | ENSMUST00000136648.7 | 1515 | No protein | lncRNA | - | - | TSL:1 |
| **Nck2-203** | ENSMUST00000124817.1 | 1421 | No protein | lncRNA | - | - | TSL:1 |

The strategy is based on the design of *Nck2-201* transcript,The transcription is shown below



# Genomic location distribution



# Protein domain



集萃药康
GemPharmatech

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

集萃药康生物科技
GemPharmatech Co.,Ltd

GemPharmatech Co.,Ltd