

# ***Birc2* Cas9-CKO Strategy**

Designer: Xueting Zhang

Design Date: 2019-8-1

# Project Overview

**Project Name**

*Birc2*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Birc2* gene. The schematic diagram is as follows:



- The *Birc2* gene has 5 transcripts. According to the structure of *Birc2* gene, exon2-exon8 of *Birc2-204* (ENSMUST00000190341.6) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Birc2* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Homozygous null mice showed a modest reduction in the number of lymphocytes. Mice homozygous for a knock-in allele exhibit increased T cell poliferation and IFNG secretion in response to anti-CD3 stimulation.
- Transcript *Birc2*-201&203 may not be affected .
- The *Birc2* gene is located on the Chr9. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Birc2 baculoviral IAP repeat-containing 2 [Mus musculus (house mouse)]

Gene ID: 11797, updated on 31-Jan-2019

### Summary




<b>Official Symbol</b>	Birc2 provided by <a href="#">MGI</a>
<b>Official Full Name</b>	baculoviral IAP repeat-containing 2 provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:1197009</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG00000057367</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	AW146227, Api1, Api2, Birc3, C-IAP1, C330006D17Rik, HIAP1, HIAP2, IAP1, IAP2, MIAP1, MIAP2, MIHB, MIHC, RNF48, ciAP1, ciAP2, mciAP1
<b>Expression</b>	Ubiquitous expression in CNS E11.5 (RPKM 9.0), CNS E14 (RPKM 8.3) and 26 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 5 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Birc2-204	<a href="#">ENSMUST00000190341.6</a>	3331	<a href="#">612aa</a>	Protein coding	<a href="#">CCDS22811</a>	<a href="#">Q62210</a>	TSL:5 GENCODE basic APPRIS P1
Birc2-202	<a href="#">ENSMUST00000074246.6</a>	3130	<a href="#">612aa</a>	Protein coding	<a href="#">CCDS22811</a>	<a href="#">Q62210</a>	TSL:1 GENCODE basic APPRIS P1
Birc2-201	<a href="#">ENSMUST00000054878.5</a>	378	<a href="#">125aa</a>	Protein coding	-	<a href="#">F6X8Q0</a>	TSL:NA GENCODE basic
Birc2-205	<a href="#">ENSMUST00000191248.1</a>	707	No protein	Processed transcript	-	-	TSL:3
Birc2-203	<a href="#">ENSMUST00000189935.1</a>	1316	No protein	Retained intron	-	-	TSL:NA

The strategy is based on the design of *Birc2-204* transcript,The transcription is shown below




# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Homozygous null mice showed a modest reduction in the number of lymphocytes. Mice homozygous for a knock-in allele exhibit increased T cell proliferation and IFNG secretion in response to anti-CD3 stimulation.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

