

# ***Fbxo34*** Cas9-CKO Strategy

Designer: Xueting Zhang

Design Date: 2019-8-2

# Project Overview

**Project Name**

***Fbxo34***

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Fbxo34* gene. The schematic diagram is as follows:



# Technical routes

- The *Fbxo34* gene has 14 transcripts. According to the structure of *Fbxo34* gene, exon2 of *Fbxo34-201* (ENSMUST00000043112.8) transcript is recommended as the knockout region. The region contains most of the coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Fbxo34* gene. The brief process is as follows: gRNA was transcribed in vitro, donor was constructed. Cas9, gRNA and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

# Notice

- Transcript *Fbxo34*-208 may not be affected.
- The *Fbxo34* gene is located on the Chr14. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Fbxo34 F-box protein 34 [Mus musculus (house mouse)]

Gene ID: 78938, updated on 31-Jan-2019

### Summary




<b>Official Symbol</b>	Fbxo34 provided by <a href="#">MGI</a>
<b>Official Full Name</b>	F-box protein 34 provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:1926188</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG000000037536</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	2900057B08Rik, 5830426G16Rik
<b>Expression</b>	Ubiquitous expression in testis adult (RPKM 13.2), liver E14.5 (RPKM 10.2) and 28 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 14 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Fbxo34-202	<a href="#">ENSMUST00000095941.8</a>	3418	<a href="#">695aa</a>	Protein coding	<a href="#">CCDS49471</a>	<a href="#">Q80XI1</a>	TSL:1 GENCODE basic APPRIS ALT2
Fbxo34-204	<a href="#">ENSMUST00000165714.1</a>	3159	<a href="#">695aa</a>	Protein coding	<a href="#">CCDS49471</a>	<a href="#">Q80XI1</a>	TSL:1 GENCODE basic APPRIS ALT2
Fbxo34-203	<a href="#">ENSMUST00000163324.7</a>	3132	<a href="#">695aa</a>	Protein coding	<a href="#">CCDS49471</a>	<a href="#">Q80XI1</a>	TSL:2 GENCODE basic APPRIS ALT2
Fbxo34-205	<a href="#">ENSMUST00000168833.8</a>	2993	<a href="#">695aa</a>	Protein coding	<a href="#">CCDS49471</a>	<a href="#">Q80XI1</a>	TSL:1 GENCODE basic APPRIS ALT2
Fbxo34-201	<a href="#">ENSMUST00000043112.8</a>	2937	<a href="#">746aa</a>	Protein coding	<a href="#">CCDS26988</a>	<a href="#">E9QM55</a>	TSL:1 GENCODE basic APPRIS P3
Fbxo34-213	<a href="#">ENSMUST00000228668.1</a>	664	<a href="#">108aa</a>	Protein coding	-	<a href="#">A0A2I3BQA4</a>	CDS 3' incomplete
Fbxo34-206	<a href="#">ENSMUST00000226395.1</a>	613	<a href="#">88aa</a>	Protein coding	-	<a href="#">A0A2I3BQU0</a>	CDS 3' incomplete
Fbxo34-207	<a href="#">ENSMUST00000226432.1</a>	594	<a href="#">93aa</a>	Protein coding	-	<a href="#">A0A2I3BRP7</a>	CDS 3' incomplete
Fbxo34-214	<a href="#">ENSMUST00000228740.1</a>	354	<a href="#">28aa</a>	Protein coding	-	<a href="#">A0A2I3BRE0</a>	CDS 3' incomplete
Fbxo34-211	<a href="#">ENSMUST00000228019.1</a>	3490	<a href="#">45aa</a>	Nonsense mediated decay	-	<a href="#">A0A2I3BPD7</a>	CDS 5' incomplete
Fbxo34-208	<a href="#">ENSMUST00000226954.1</a>	505	<a href="#">51aa</a>	Nonsense mediated decay	-	<a href="#">A0A2I3BR79</a>	CDS 5' incomplete
Fbxo34-210	<a href="#">ENSMUST00000227601.1</a>	393	<a href="#">44aa</a>	Nonsense mediated decay	-	<a href="#">A0A2I3BS05</a>	CDS 5' incomplete
Fbxo34-209	<a href="#">ENSMUST00000227162.1</a>	373	No protein	Processed transcript	-	-	
Fbxo34-212	<a href="#">ENSMUST00000228634.1</a>	338	No protein	Retained intron	-	-	

The strategy is based on the design of *Fbxo34-201* transcript,The transcription is shown below



# Genomic location distribution





# Protein domain



If you have any questions, you are welcome to inquire.

Tel: 400-9660890

