

# ONDHOM DECK Donal Day Co. Il27 Cas9-CKO Strategy Rohalmakech Co.

Conposition of the Constitution of the Constit

Conday are ch JiaYu

# **Project Overview**



**Project Name** 

*Il27* 

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Il27* gene. The schematic diagram is as follows:



## Technical routes



- ➤ The *Il27* gene has 1 transcript. According to the structure of *Il27* gene, exon2-exon3 of *Il27-201*(ENSMUST0000058429.5) transcript is recommended as the knockout region. The region contains 260bp coding sequence.

  Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Il27* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

## **Notice**



- ➤ According to the existing MGI data, Mice homozygous for a knock-out allele develop an exaggerated delayed-type hypersensitivity response and are more susceptible to experimental autoimmune encephalomyelitis due to the presence of an increased number of Th17 cells.
- > The *Il27* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

## Gene information (NCBI)



#### II27 interleukin 27 [Mus musculus (house mouse)]

Gene ID: 246779, updated on 12-Mar-2019

#### Summary

☆ ?

Official Symbol II27 provided by MGI

Official Full Name interleukin 27 provided by MGI

Primary source MGI:MGI:2384409

See related Ensembl: ENSMUSG00000044701

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as IL-27, IL-27-A, IL-27p28, IL27-A, II30, p28

Expression Broad expression in genital fat pad adult (RPKM 5.8), spleen adult (RPKM 2.5) and 18 other tissuesSee more

Orthologs <u>human</u> all


# Transcript information (Ensembl)



The gene has 1 transcript, and the transcript is shown below:


| Name     | Transcript ID        | bp   | Protein | Biotype        | CCDS      | UniProt | Flags                         |   |
|----------|----------------------|------|---------|----------------|-----------|---------|-------------------------------|---|
| 1127-201 | ENSMUST00000058429.5 | 1021 | 234aa   | Protein coding | CCDS21835 | Q8K3I6  | TSL:1 GENCODE basic APPRIS P1 | ľ |

The strategy is based on the design of *Il27-201* transcript, The transcription is shown below




## Genomic location distribution






## Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Mice homozygous for a knock-out allele develop an exaggerated delayed-type hypersensitivity response and are more susceptible to experimental autoimmune encephalomyelitis due to the presence of an increased number of Th17 cells.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





