

Kdm8 Cas9-CKO Strategy

Designer: Xiaojing Li

Design Date: 2019-9-11

Reviewer: JiaYu

Project Overview



Project Name

Kdm8

Project type

Cas9-CKO


Strain background

C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Kdm8* gene. The schematic diagram is as follows:



Technical routes



- The *Kdm8* gene has 2 transcripts. According to the structure of *Kdm8* gene, exon3 of *Kdm8-201*(ENSMUST00000033010.8) transcript is recommended as the knockout region. The region contains 167bp coding sequence.

 Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Kdm8* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- ➤ According to the existing MGI data, Homozygous inactivation of this gene leads to complete embryonic lethality during organogenesis associated with severe growth retardation and abnormal embryo turning. Observed phenotypes include open neural tubes and absent vitelline blood vessels.
- > The *Kdm8* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Kdm8 lysine (K)-specific demethylase 8 [Mus musculus (house mouse)]

Gene ID: 77035, updated on 19-Mar-2019

Summary

☆ ?

Official Symbol Kdm8 provided by MGI

Official Full Name lysine (K)-specific demethylase 8 provided by MGI

Primary source MGI:MGI:1924285

See related Ensembl: ENSMUSG00000030752

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as 3110005O21Rik, Jmjd5

Expression Ubiquitous expression in thymus adult (RPKM 9.3), ovary adult (RPKM 7.0) and 28 other tissuesSee more

Orthologs <u>human</u> all


Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Kdm8-201	ENSMUST00000033010.8	2389	414aa	Protein coding	CCDS21821	Q9CXT6	TSL:1 GENCODE basic APPRIS P1
Kdm8-202	ENSMUST00000135129.1	601	<u>152aa</u>	Protein coding	-8	D3YUV8	CDS 3' incomplete TSL:3

The strategy is based on the design of *Kdm8-201* transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous inactivation of this gene leads to complete embryonic lethality during organogenesis associated with severe growth retardation and abnormal embryo turning. Observed phenotypes include open neural tubes and absent vitelline blood vessels.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





