

# Olfr95 Cas9-KO Strategy

**Designer:** 

Reviewer

**Design Date:** 

Ruirui Zhang

**Huimin Su** 

2019-8-16

## **Project Overview**



**Project Name** 

Olfr95

**Project type** 

Cas9-KO


Strain background

C57BL/6JGpt

# **Knockout strategy**



This model will use CRISPR/Cas9 technology to edit the Olfr95 gene. The schematic diagram is as follows:



### **Technical routes**



- ➤ The *Olfr95* gene has 2 transcripts. According to the structure of *Olfr95* gene, exon1 of *Olfr95-201* (ENSMUST00000060728.6) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Olfr95* gene. The brief process is as follows: CRISPR/Cas9 system

### **Notice**



- > The *Olfr95* gene is located on the Chr17. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- ➤ Olfr96 gene is about 7.0kb away from Olfr95 gene. Olfr95 gene knockout may affect the regulation of Olfr96 gene at 5 terminals.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

### Gene information (NCBI)



#### Olfr95 olfactory receptor 95 [ Mus musculus (house mouse) ]

Gene ID: 258506, updated on 12-Aug-2019

#### Summary

☆ ?

Official Symbol Olfr95 provided by MGI

Official Full Name olfactory receptor 95 provided by MGI

Primary source MGI:MGI:2177478

See related Ensembl: ENSMUSG00000049561

Gene type protein coding
RefSeq status PROVISIONAL
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae;

Murinae; Mus; Mus

Also known as MOR263-6

Summary Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor

proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes

and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008]

Orthologs human all


# Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Olfr95-201	ENSMUST00000060728.6	5129	312aa	Protein coding	CCDS28739	Q8VFE2	TSL:NA GENCODE basic APPRIS P1
Olfr95-202	ENSMUST00000216318.1	4917	312aa	Protein coding	CCDS28739	Q8VFE2	TSL:5 GENCODE basic APPRIS P1

The strategy is based on the design of Olfr95-201 transcript, The transcription is shown below




### Genomic location distribution





### Protein domain







If you have any questions, you are welcome to inquire. Tel: 400-9660890





