

Cenpa Cas9-CKO Strategy

Designer:Xiaojing Li

Design Date:2019-11-18

Reviewer:Jia Yu

Project Overview



Project Name Cenpa


Project type Cas9-CKO

Strain background C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the Cenpa gene. The schematic diagram is as follows:



Technical routes



- The *Cenpa* gene has 10 transcripts. According to the structure of *Cenpa* gene, exon1 of *Cenpa-205*(ENSMUST00000144742.5) transcript is recommended as the knockout region. The region contains start codon ATG.

 Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cenpa* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- > According to the existing MGI data, Homozygous mutation of this gene results in embryonic lethality between E3.5 and E10.5. Embryogenesis is impaired due to chromosomal missegregation, aneuploidy, and apoptosis.
- The *Cenpa* gene is located on the Chr5. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Cenpa centromere protein A [Mus musculus (house mouse)]

Gene ID: 12615, updated on 10-Oct-2019



☆ ?

Official Symbol Cenpa provided by MGI

Official Full Name centromere protein A provided by MGI

Primary source MGI:MGI:88375

See related Ensembl: ENSMUSG00000029177

Gene type protein coding
RefSeq status REVIEWED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as Cenp-A

Summary Centromeres are the differentiated chromosomal domains that specify the mitotic behavior of chromosomes. This gene encodes a

centromere protein which contains a histone H3 related histone fold domain that is required for targeting to the centromere. Centromere protein A is proposed to be a component of a modified nucleosome or nucleosome-like structure in which it replaces 1 or both copies of conventional histone H3 in the (H3-H4)2 tetrameric core of the nucleosome particle. The protein is a replication-independent histone that is a member of the histone H3 family. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by

RefSeq. Nov 2015]

Expression Broad expression in CNS E11.5 (RPKM 43.5), liver E14.5 (RPKM 41.9) and 20 other tissues See more

Orthologs human all


Transcript information (Ensembl)



The gene has 10 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Cenpa-205	ENSMUST00000144742.5	1466	<u>134aa</u>	Protein coding	CCDS19162	<u>O35216</u>	TSL:1 GENCODE basic APPRIS P1
Cenpa-209	ENSMUST00000199320.4	369	<u>99aa</u>	Protein coding	*	A0A0G2JGI2	CDS 3' incomplete TSL:3
Cenpa-206	ENSMUST00000149759.1	346	<u>59aa</u>	Protein coding	-	A0A0G2JEV0	TSL:3 GENCODE basic
Cenpa-202	ENSMUST00000133316.7	1031	<u>42aa</u>	Nonsense mediated decay		D6RCV6	TSL:1
Cenpa-204	ENSMUST00000134846.7	1007	<u>39aa</u>	Nonsense mediated decay	-	D6RJ71	TSL:2
Cenpa-201	ENSMUST00000031073.9	993	<u>39aa</u>	Nonsense mediated decay	-8	D6RJ71	TSL:1
Cenpa-210	ENSMUST00000199617.1	827	<u>130aa</u>	Nonsense mediated decay	-	A0A0G2JEV2	TSL:3
Cenpa-208	ENSMUST00000197962.1	3806	No protein	Retained intron		10	TSL:NA
Cenpa-207	ENSMUST00000150810.7	1120	No protein	Retained intron	-	9	TSL:5
Cenpa-203	ENSMUST00000134372.1	1010	No protein	Retained intron	-8		TSL:2

The strategy is based on the design of Cenpa-205 transcript, The transcription is shown below



Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous mutation of this gene results in embryonic lethality between E3.5 and E10.5. Embryogenesis is impaired due to chromosomal missegregation, aneuploidy, and apoptosis.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





