

Cyp7a1 Cas9-KO Strategy

Designer:

Daohua Xu

Reviewer:

Huimin Su

Design Date:

2019-11-14

Project Overview

Project Name

Cyp7a1

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Cyp7a1* gene. The schematic diagram is as follows:



- The *Cyp7a1* gene has 2 transcripts. According to the structure of *Cyp7a1* gene, exon2-exon4 of *Cyp7a1-201* (ENSMUST00000029905.1) transcript is recommended as the knockout region. The region contains 959bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cyp7a1* gene. The brief process is as follows: CRISPR/Cas9 system

- According to the existing MGI data, Mice homozygous for disruption of this gene experience severe neonatal and postnatal lethality. Supplementation of the maternal diet with fat soluble vitamins and cholic acid starting before birth alleviates much of the phenotype.
- The *Cyp7a1* gene is located on the Chr4. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Cyp7a1 cytochrome P450, family 7, subfamily a, polypeptide 1 [Mus musculus (house mouse)]

Gene ID: 13122, updated on 19-Mar-2019

Summary




Official Symbol	Cyp7a1 provided by MGI
Official Full Name	cytochrome P450, family 7, subfamily a, polypeptide 1 provided by MGI
Primary source	MGI:MGI:106091
See related	Ensembl:ENSMUSG000000028240
Gene type	protein coding
RefSeq status	VALIDATED
Organism	Mus musculus
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	CYPVII, CYPVIIc
Expression	Biased expression in liver E18 (RPKM 9.1) and liver adult (RPKM 7.9) See more
Orthologs	human all

Transcript information (Ensembl)


The gene has 2 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Cyp7a1-201	ENSMUST00000029905.1	4172	503aa	Protein coding	CCDS17950	Q64505	TSL:1 GENCODE basic APPRIS P1
Cyp7a1-202	ENSMUST00000147346.1	228	No protein	lncRNA	-	-	TSL:5


The strategy is based on the design of *Cyp7a1-201* transcript,The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous for disruption of this gene experience severe neonatal and postnatal lethality. Supplementation of the maternal diet with fat soluble vitamins and cholic acid starting before birth alleviates much of the phenotype.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

