

S1pr2 Cas9-CKO Strategy

Designer:

Daohua Xu

Reviewer:

Huimin Su

Design Date:

2019-9-10

Project Overview



Project Name

S1pr2

Project type

Cas9-CKO


Strain background

C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the S1pr2 gene. The schematic diagram is as follows:



Technical routes



- The *S1pr2* gene has 4 transcripts. According to the structure of *S1pr2* gene, exon2 of *S1pr2-201* (ENSMUST00000054197.6) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *S1pr2* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- > The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- > According to the existing MGI data, Homozygous null mutations in this gene may lead to impaired auditory and vestibular function, multiple inner ear pathologies, deafness, altered neuronal excitability, lethal seizures, altered physiology of germinal center B cells, small litter size, and enhanced tumor angiogenesis and tumor growth.
- ➤ This strategy may affect the 5-terminal regulation of the *Dnmt1* gene.
- > The *S1pr2* gene is located on the Chr9. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



S1pr2 sphingosine-1-phosphate receptor 2 [Mus musculus (house mouse)]

Gene ID: 14739, updated on 12-Mar-2019

Summary

☆ ?

Official Symbol S1pr2 provided by MGI

Official Full Name sphingosine-1-phosphate receptor 2 provided by MGI

Primary source MGI:MGI:99569

See related Ensembl: ENSMUSG00000043895

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as 1100001A16Rik, Edg5, Gpcr13, H218, LPb2, S1P2

Expression Broad expression in lung adult (RPKM 19.6), limb E14.5 (RPKM 17.0) and 24 other tissuesSee more

Orthologs <u>human</u> all

Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
S1pr2-201	ENSMUST00000054197.6	6394	352aa	Protein coding	CCDS22888	P52592	TSL:1 GENCODE basic APPRIS P1
S1pr2-202	ENSMUST00000214218.1	379	<u>16aa</u>	Protein coding	-	A0A1L1ST05	CDS 3' incomplete TSL:3
S1pr2-204	ENSMUST00000216499.1	419	No protein	Processed transcript	-	-	TSL:2
S1pr2-203	ENSMUST00000214333.1	1587	No protein	Retained intron	22	12	TSL:NA


The strategy is based on the design of S1pr2-201 transcript, The transcription is shown below

< S1pr2-201
protein coding

Reverse strand — 14.42 kb —


Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous null mutations in this gene may lead to impaired auditory and vestibular function, multiple inner ear pathologies, deafness, altered neuronal excitability, lethal seizures, altered physiology of germinal center B cells, small litter size, and enhanced tumor angiogenesis and tumor growth.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





