

# Slc15a3 Cas9-CKO Strategy Rohalanakoch Co.

JiaYu

# **Project Overview**



**Project Name** 

Slc15a3

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the Slc15a3 gene. The schematic diagram is as follows:



## Technical routes



- ➤ The *Slc15a3* gene has 4 transcripts. According to the structure of *Slc15a3* gene, exon2 of *Slc15a3-201*(ENSMUST00000025646.2) transcript is recommended as the knockout region. The region contains 284bp coding sequence.

  Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Slc15a3* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

## **Notice**



- > According to the existing MGI data, The gene is involved in pathogen sensing by dendritic cells.

  Homozygous KO results in a reduction of the number of these cells displaying tubular endo-lysosomes after LPS treatment.
- The *Slc15a3* gene is located on the Chr19. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

## Gene information (NCBI)



#### Slc15a3 solute carrier family 15, member 3 [Mus musculus (house mouse)]

Gene ID: 65221, updated on 31-Jan-2019

#### Summary

☆ ?

Official Symbol Slc15a3 provided by MGI

Official Full Name solute carrier family 15, member 3 provided by MGI

Primary source MGI:MGI:1929691

See related Ensembl: ENSMUSG00000024737

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as Ci1, cl-1

Expression Broad expression in spleen adult (RPKM 18.7), lung adult (RPKM 15.6) and 17 other tissuesSee more

Orthologs <u>human</u> all

# Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
SIc15a3-201	ENSMUST00000025646.2	2345	<u>578aa</u>	Protein coding	CCDS29588	Q8BPX9	TSL:1 GENCODE basic APPRIS P1
SIc15a3-204	ENSMUST00000138380.1	414	No protein	Processed transcript	*	0.00	TSL:2
SIc15a3-202	ENSMUST00000131010.1	361	No protein	Processed transcript	S-	323	TSL:3
SIc15a3-203	ENSMUST00000138263.1	4868	No protein	Retained intron	92	127	TSL:1

The strategy is based on the design of Slc15a3-201 transcript, The transcription is shown below




## Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, The gene is involved in pathogen sensing by dendritic cells. Homozygous KO results in a reduction of the number of these cells displaying tubular endo-lysosomes after LPS treatment.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





