

# Donald Color Sf3b1 Cas9-KO Strategy The state of the s Conposition of the contraction o

Condand Stock JiaYu

# **Project Overview**



**Project Name** 

Sf3b1

**Project type** 

Cas9-KO


Strain background

C57BL/6JGpt

# **Knockout strategy**



This model will use CRISPR/Cas9 technology to edit the Sf3b1 gene. The schematic diagram is as follows:



## **Technical routes**



- ➤ The *Sf3b1* gene has 8 transcripts. According to the structure of *Sf3b1* gene, exon2-exon5 of *Sf3b1-201*(ENSMUST00000027127.13) transcript is recommended as the knockout region. The region contains 467bp coding sequence Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify Sf3b1 gene. The brief process is as follows: CRISPR/Cas9 system

### **Notice**



- ➤ According to the existing MGI data, Homozygous null embryos die around the 16- to 32-cell stage. Heterozygous mice exhibit various skeletal transformations.
- > The *Sf3b1* gene is located on the Chr1. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- > This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

## Gene information (NCBI)



#### Sf3b1 splicing factor 3b, subunit 1 [Mus musculus (house mouse)]

Gene ID: 81898, updated on 26-Feb-2019

#### Summary

☆ ?

Official Symbol Sf3b1 provided by MGI

Official Full Name splicing factor 3b, subunit 1 provided by MGI

Primary source MGI:MGI:1932339

See related Ensembl:ENSMUSG00000025982

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as 155kDa, 2810001M05Rik, AA409119, Prp10, SAP155, SF3b155, TA-8, Targ4

Expression Ubiquitous expression in CNS E14 (RPKM 50.4), CNS E11.5 (RPKM 49.8) and 25 other tissuesSee more

Orthologs <u>human</u> all


# Transcript information (Ensembl)



The gene has 8 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Sf3b1-201	ENSMUST00000027127.13	6195	<u>1304aa</u>	Protein coding	CCDS35567	G5E866	TSL:1 GENCODE basic APPRIS P1
Sf3b1-208	ENSMUST00000191303.1	600	<u>144aa</u>	Nonsense mediated decay		A0A087WNS2	TSL:5
Sf3b1-206	ENSMUST00000189051.1	804	No protein	Processed transcript	-	40	TSL:3
Sf3b1-202	ENSMUST00000185429.1	3589	No protein	Retained intron	-	20	TSL:NA
Sf3b1-203	ENSMUST00000187500.6	3403	No protein	Retained intron		T4	TSL:2
Sf3b1-207	ENSMUST00000190175.1	3350	No protein	Retained intron		<del>.</del>	TSL:2
Sf3b1-204	ENSMUST00000188419.1	711	No protein	Retained intron	-	40	TSL:2
Sf3b1-205	ENSMUST00000188859.6	628	No protein	Retained intron	- <u>-</u>	29	TSL:2

The strategy is based on the design of Sf3b1-201 transcript, The transcription is shown below




## Genomic location distribution






## Protein domain





## Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous null embryos die around the 16- to 32-cell stage. Heterozygous mice exhibit various skeletal transformations.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





