

# Id1 Cas9-CKO Strategy

Designer: Daohua Xu

Reviewer: Huimin Su

**Design Date:** 2019-11-26

# **Project Overview**



**Project Name** 

Id1

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Id1* gene. The schematic diagram is as follows:



### Technical routes



- The *Id1* gene has 2 transcripts. According to the structure of *Id1* gene, exon1-exon2 of *Id1-201* (ENSMUST00000038368.8) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Id1* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

### **Notice**



- ➤ According to the existing MGI data, Homozygotes for knockout alleles of both Id1 and Id3 exhibit vascular malformations in the forebrain, lack of vascular branching and sprouting in the neuroectoderm, and impaired angiogenesis in transplanted and spontaneous tumors.
- > The *Id1* gene is located on the Chr2. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

### Gene information (NCBI)



#### Id1 inhibitor of DNA binding 1, HLH protein [Mus musculus (house mouse)]

Gene ID: 15901, updated on 19-Mar-2019

#### Summary

^ ?

Official Symbol Id1 provided by MGI

Official Full Name inhibitor of DNA binding 1, HLH protein provided by MGI

Primary source MGI:MGI:96396

See related Ensembl: ENSMUSG00000042745

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as Al323524, D2Wsu140e, ldb1, bHLHb24

Expression Broad expression in stomach adult (RPKM 416.7), colon adult (RPKM 399.7) and 22 other tissuesSee more

Orthologs human all

# Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
ld1-201	ENSMUST00000038368.8	934	148aa	Protein coding	CCDS16897	P20067 Q6GTZ3	TSL:1 GENCODE basic APPRIS P2
ld1-202	ENSMUST00000109824.1	1160	<u>168aa</u>	Protein coding	( <del>-</del> 8	A2AHY3	TSL:NA GENCODE basic APPRIS ALT2

The strategy is based on the design of *Id1-201* transcript, The transcription is shown below




### Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygotes for knockout alleles of both Id1 and Id3 exhibit vascular malformations in the forebrain, lack of vascular branching and sprouting in the neuroectoderm, and impaired angiogenesis in transplanted and spontaneous tumors.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





