

# *Slc5a5* Cas9-CKO Strategy

Designer: Yanhua Shen

# Project Overview

**Project Name**

*Slc5a5*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Slc5a5* gene. The schematic diagram is as follows:



- The *Slc5a5* gene has 1 transcript. According to the structure of *Slc5a5* gene, exon2-exon13 of *Slc5a5-201* (ENSMUST00000000809.2) transcript is recommended as the knockout region. The region contains 1279bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Slc5a5* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

# Notice

- According to the existing MGI data, Mice homozygous for a knock-out allele exhibit reduced T3 and T4 levels when fed a minimal iodine diet.
- Intron1-2 is 587 bp, the insertion of loxp may affect the splicing of this gene.
- The N-terminus may remain around 120 amino acids, and some functions may be retained.
- The *Slc5a5* gene is located on the Chr8. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Slc5a5 solute carrier family 5 (sodium iodide symporter), member 5 [Mus musculus (house mouse)]

Gene ID: 114479, updated on 31-Jan-2019

### Summary




<b>Official Symbol</b>	Slc5a5 provided by <a href="#">MGI</a>
<b>Official Full Name</b>	solute carrier family 5 (sodium iodide symporter), member 5 provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:2149330</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG000000000792</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	NIS
<b>Expression</b>	Biased expression in stomach adult (RPKM 130.4) and mammary gland adult (RPKM 64.6) <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 1 transcript, and the transcript is shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Slc5a5-201	<a href="#">ENSMUST00000000809.2</a>	2928	<a href="#">618aa</a>	Protein coding	<a href="#">CCDS22385</a>	<a href="#">G3X8P5</a>	TSL:1 GENCODE basic APPRIS P1

The strategy is based on the design of *Slc5a5-201* transcript, The transcription is shown below




# Genomic location distribution





# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Mice homozygous for a knock-out allele exhibit reduced T3 and T4 levels when fed a minimal iodine diet.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

