

# Slc39a10 Cas9-CKO Strategy

Designer: Yanhua Shen

Reviewer:Xueting Zhang

Design Date:2019-08-30

# **Project Overview**



**Project Name** 

Slc39a10

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

# Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the Slc39a10 gene. The schematic diagram is as follows:



### Technical routes



- The *Slc39a10* gene has 4 transcripts. According to the structure of *Slc39a10* gene, exon2-exon9 of *Slc39a10-201* (ENSMUST00000027131.5) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Slc39a10* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

### **Notice**



- > According to the existing MGI data, Mice with conditional loss of function display defects in cellular proliferation and differentiation.
- ➤ The effects of transcripts 203 and 204 is unknown.
- The *Slc39a10* gene is located on the Chr1. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- ➤ This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

### Gene information (NCBI)



#### Slc39a10 solute carrier family 39 (zinc transporter), member 10 [ Mus musculus (house mouse) ]

Gene ID: 227059, updated on 14-Aug-2019

#### Summary

☆ ?

Official Symbol Slc39a10 provided by MGI

Official Full Name solute carrier family 39 (zinc transporter), member 10 provided by MGI

Primary source MGI:MGI:1914515

See related Ensembl: ENSMUSG00000025986

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as Zip10; mKIAA1265; 5430433I10; 2900042E17Rik

Expression Broad expression in cortex adult (RPKM 17.1), frontal lobe adult (RPKM 14.5) and 22 other tissues See more

Orthologs human all


# Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:


Name 🍦	Transcript ID 👙	bp 🌲	Protein 4	Biotype	CCDS 🍦	UniProt 🍦	Flags
Slc39a10-201	ENSMUST00000027131.5	5498	<u>833aa</u>	Protein coding	CCDS14936₽	Q6P5F6₽	TSL:1 GENCODE basic APPRIS P1
Slc39a10-204	ENSMUST00000186852.1	565	<u>167aa</u>	Protein coding	14	A0A087WQF7₽	CDS 3' incomplete TSL:3
Slc39a10-203	ENSMUST00000185520.1	497	<u>144aa</u>	Protein coding	14	<u>A0A087WRC8</u> €	CDS 3' incomplete TSL:3
Slc39a10-202	ENSMUST00000141226.2	2372	No protein	Retained intron	12	640	TSL:1

The strategy is based on the design of Slc39a10-201 transcript, The transcription is shown below




### Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Mice with conditional loss of function display defects in cellular proliferation and differentiation.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





