

Trpc5 Cas9-CKO Strategy

Designer:

Yang Zeng

Reviewer:

Xueting Zhang

Design Date:

2019-10-28

Project Overview

Project Name

Trpc5

Project type


Cas9-CKO

Strain background

C57BL/6JGpt

Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Trpc5* gene. The schematic diagram is as follows:



- The *Trpc5* gene has 2 transcripts. According to the structure of *Trpc5* gene, exon4-exon6 of *Trpc5-201* (ENSMUST00000040184.3) transcript is recommended as the knockout region. The region contains 800bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Trpc5* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Mice with homozygous null alleles of the gene exhibit diminished innate fear response with reduction in synaptic activation and strength in neurons of the amygdala.
- The N-terminal of *Trpc5* gene will remain several amino acids, it may remain the partial function of *Trpc5* gene.
- The *Trpc5* gene is located on the ChrX. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.


Gene information (NCBI)

Trpc5 transient receptor potential cation channel, subfamily C, member 5 [*Mus musculus* (house mouse)]

Gene ID: 22067, updated on 10-Sep-2019

Summary

Official Symbol Trpc5 provided by MGI
Official Full Name transient receptor potential cation channel, subfamily C, member 5 provided by MGI
Primary source [MGI:MGI:109524](#)
See related [Ensembl:ENSMUSG00000041710](#)
Gene type protein coding
RefSeq status VALIDATED
Organism [Mus musculus](#)
Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as CCE2; TRP5; TRP-5; Trrp5
Expression Biased expression in CNS E18 (RPKM 1.2), frontal lobe adult (RPKM 1.0) and 6 other tissues [See more](#)
Orthologs [human](#) [all](#)




Transcript information (Ensembl)


The gene has 2 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Translation ID	Biotype	CCDS	UniProt	Flags
Trpc5-201	ENSMUST00000040184.3	4158	975aa	ENSMUSP00000049063.3	Protein coding	CCDS30457	Q2KHN9 Q9QX29	TSL:1 Gencode basic APPRIS P1
Trpc5-202	ENSMUST00000148240.1	2782	No protein	-	Retained intron	-	-	TSL:1


The strategy is based on the design of *Trpc5-201* transcript,The transcription is shown below



Genomic location distribution




Protein domain



Mouse phenotype description(MGI)

Phenotype Overview ?



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice with homozygous null alleles of the gene exhibit diminished innate fear response with reduction in synaptic activation and strength in neurons of the amygdala.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

