

# Kcnk3 Cas9-CKO Strategy

Designer:Xueting Zhang

Reviewer: Yanhua Shen

Date:2019-11-24

# **Project Overview**



**Project Name** 

Kcnk3

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Kcnk3* gene. The schematic diagram is as follows:



### **Technical routes**



- ➤ The *Kcnk3* gene has 2 transcripts. According to the structure of *Kcnk3* gene, exon2 of *Kcnk3-201*(ENSMUST00000066295.3) transcript is recommended as the knockout region. The region contains most of the coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Kcnk3* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- > The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

### **Notice**



- ➤ According to the existing MGI data, Mice homozygous for a null alleles exhibit decreased pH sensitivitive of action potential in serotonergic neurons.
- > The *Kcnk3* gene is located on the Chr5. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

### Gene information (NCBI)



#### Kcnk3 potassium channel, subfamily K, member 3 [ Mus musculus (house mouse) ]

Gene ID: 16527, updated on 12-Aug-2019

#### Summary

☆ ?

Official Symbol Kcnk3 provided by MGI

Official Full Name potassium channel, subfamily K, member 3 provided by MGI

Primary source MGI:MGI:1100509

See related Ensembl: ENSMUSG00000049265

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea;

Muridae; Murinae; Mus; Mus

Also known as TASK; Task-1; cTBAK-1

Expression Biased expression in adrenal adult (RPKM 1700.1) and heart adult (RPKM 66.4) See more

Orthologs human all

#### **Genomic context**



**Location:** 5 B1; 5 16.68 cM

See Kcnk3 in Genome Data Viewer

Exon count: 2

Annotation release	Status	Assembly	Chr	Location	
108	current	GRCm38.p6 (GCF_000001635.26)	5	NC_000071.6 (3058817030625270)	
Build 37.2	previous assembly	MGSCv37 (GCF 000001635.18)	5	NC 000071.5 (3089054330927643)	

# Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Kcnk3-201	ENSMUST00000066295.3	3811	<u>409aa</u>	Protein coding	CCDS19160	<u>035111</u>	TSL:1 GENCODE basic APPRIS P1
Kcnk3-202	ENSMUST00000197428.1	507	No protein	IncRNA	-	(#1	TSL:2

The strategy is based on the design of Kcnk3-201 transcript, The transcription is shown below

Kcnk3-201 > protein coding


### Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Mice homozygous for a null alleles exhibit decreased pH sensitivitive of action potential in serotonergic neurons.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





