

# ***Txk* Cas9-CKO Strategy**

**Designer:**

**Daohua Xu**

**Reviewer:**

**Huimin Su**

**Design Date:**

**2020-2-12**

# Project Overview

---

**Project Name**

***Txk***

---

**Project type**

**Cas9-CKO**

---


**Strain background**

**C57BL/6JGpt**

---

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Txk* gene. The schematic diagram is as follows:



- The *Txk* gene has 8 transcripts. According to the structure of *Txk* gene, exon2-exon5 of *Txk*-202 (ENSMUST00000169534.5) transcript is recommended as the knockout region. The region contains 427bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Txk* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Homozygous mutation of this gene results in increased susceptibility to parasitic (*Toxoplasma gondii*) infection and decreased cytokine secretion in stimulated splenocytes.
- The KO region contains functional region of the *Gm43717* gene. Knockout the region will affect the function of *Gm43717* gene.
- The *Txk* gene is located on the Chr5. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Txk TXK tyrosine kinase [Mus musculus (house mouse)]

Gene ID: 22165, updated on 31-Jan-2019

### Summary




<b>Official Symbol</b>	Txk provided by <a href="#">MGI</a>
<b>Official Full Name</b>	TXK tyrosine kinase provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:102960</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG00000054892</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	A130089B16Rik, Btkl, PTK-RL-18, PTK4, Rik
<b>Expression</b>	Biased expression in thymus adult (RPKM 5.4), spleen adult (RPKM 1.6) and 3 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 8 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
<b>Txk-202</b>	<a href="#">ENSMUST00000169534.5</a>	2344	<a href="#">527aa</a>	Protein coding	<a href="#">CCDS19334</a>	<a href="#">B2RQ20</a>	TSL:1 GENCODE basic APPRIS P3
<b>Txk-206</b>	<a href="#">ENSMUST00000198464.2</a>	2271	<a href="#">473aa</a>	Protein coding	<a href="#">CCDS80298</a>	<a href="#">P42682</a>	TSL:1 GENCODE basic
<b>Txk-201</b>	<a href="#">ENSMUST00000113604.9</a>	2260	<a href="#">527aa</a>	Protein coding	<a href="#">CCDS51514</a>	<a href="#">P42682</a>	TSL:1 GENCODE basic APPRIS ALT2
<b>Txk-203</b>	<a href="#">ENSMUST00000197313.4</a>	2190	<a href="#">505aa</a>	Protein coding	<a href="#">CCDS80299</a>	<a href="#">A0A0G2JG94</a>	TSL:1 GENCODE basic
<b>Txk-204</b>	<a href="#">ENSMUST00000197843.2</a>	5303	No protein	Retained intron	-	-	TSL:2
<b>Txk-205</b>	<a href="#">ENSMUST00000197923.2</a>	4210	No protein	Retained intron	-	-	TSL:NA
<b>Txk-208</b>	<a href="#">ENSMUST00000198970.4</a>	2439	No protein	lncRNA	-	-	TSL:1
<b>Txk-207</b>	<a href="#">ENSMUST00000198798.4</a>	2276	No protein	lncRNA	-	-	TSL:1

The strategy is based on the design of *Txk-202* transcript,The transcription is shown below




# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Homozygous mutation of this gene results in increased susceptibility to parasitic (*Toxoplasma gondii*) infection and decreased cytokine secretion in stimulated splenocytes.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

