

Pstpip2 Cas9-CKO Strategy

Designer: Daohua Xu

Reviewer: Huimin Su

Design Date: 2020-2-11

Project Overview



Project Name

Pstpip2

Project type

Cas9-CKO


Strain background

C57BL/6JGpt

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Pstpip2* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Pstpip2* gene has 2 transcripts. According to the structure of *Pstpip2* gene, exon2-exon4 of *Pstpip2-201*(ENSMUST00000114741.3) transcript is recommended as the knockout region. The region contains 214bp coding sequence.

 Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Pstpip2* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- ➤ According to the existing MGI data, Homozygous mutant animals develop osteomyelitis (bone inflammation). Tail kinks are observed starting at 6-8 weeks of age and chronic inflammation of the extremities and ears is subsequently seen. Extramedullary hematopoiesis in the spleen is observed.
- ➤ The *Pstpip2* gene is located on the Chr18. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Pstpip2 proline-serine-threonine phosphatase-interacting protein 2 [Mus musculus (house mouse)]

Gene ID: 19201, updated on 31-Jan-2019

Summary

☆ ?

Official Symbol Pstpip2 provided by MGI

Official Full Name proline-serine-threonine phosphatase-interacting protein 2 provided by MGI

Primary source MGI:MGI:1335088

See related Ensembl:ENSMUSG00000025429

Gene type protein coding
RefSeq status VALIDATED

Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as MAYP, cmo

Expression Annotation category: partial on reference assembly See more


Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Pstpip2-201	ENSMUST00000114741.3	2267	<u>334aa</u>	Protein coding	CCDS37868	Q99M15	TSL:1 GENCODE basic APPRIS P1
Pstpip2-202	ENSMUST00000238172.1	1056	230aa	Protein coding	-		GENCODE basic

The strategy is based on the design of Pstpip2-201 transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous mutant animals develop osteomyelitis (bone inflammation). Tail kinks are observed starting at 6-8 weeks of age and chronic inflammation of the extremities and ears is subsequently seen. Extramedullary hematopoiesis in the spleen is observed.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





