

Erk Cas9-CKO Strategy

Designer:

JiaYu

Reviewer:

Xiaojing Li

Design Date:

2020-2-11

Project Overview

Project Name

Frk

Project type


Cas9-CKO

Strain background

C57BL/6JGpt

Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Frk* gene. The schematic diagram is as follows:



Technical routes

- The *Frk* gene has 4 transcripts. According to the structure of *Frk* gene, exon3 of *Frk-201* (ENSMUST00000019913.14) transcript is recommended as the knockout region. The region contains 122bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Frk* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Mice homozygous for a targeted null mutation do not exhibit increased susceptibility to spontaneous tumors nor increased sensitivity to ionizing radiation. Epithelial tissues appear similar to controls, but circulating levels of T3 were significantly reduced.
- The *Frk* gene is located on the Chr10. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)

Frk fyn-related kinase [Mus musculus (house mouse)]

Gene ID: 14302, updated on 31-Jan-2019

Summary




Official Symbol	Frk provided by MGI
Official Full Name	fyn-related kinase provided by MGI
Primary source	MGI:MGI:103265
See related	Ensembl:ENSMUSG00000019779
Gene type	protein coding
RefSeq status	VALIDATED
Organism	Mus musculus
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	BSK, BSK/IYK, C85044, GTK, RAK
Expression	Broad expression in large intestine adult (RPKM 4.3), placenta adult (RPKM 3.4) and 17 other tissues See more
Orthologs	human all

Transcript information (Ensembl)


The gene has 4 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Frk-201	ENSMUST00000019913.14	4696	512aa	Protein coding	CCDS23781	Q922K9	TSL:1 GENCODE basic APPRIS P1
Frk-202	ENSMUST00000170771.2	3406	512aa	Protein coding	CCDS23781	Q922K9	TSL:1 GENCODE basic APPRIS P1
Frk-204	ENSMUST00000215594.1	2038	No protein	Retained intron	-	-	TSL:1
Frk-203	ENSMUST00000215144.1	350	No protein	lncRNA	-	-	TSL:3


The strategy is based on the design of *Frk-201* transcript,The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous for a targeted null mutation do not exhibit increased susceptibility to spontaneous tumors nor increased sensitivity to ionizing radiation. Epithelial tissues appear similar to controls, but circulating levels of T3 were significantly reduced.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

