

# Dusp5 Cas9-CKO Strategy

Designer: Huimin Su

Reviewer: Ruirui Zhang

**Design Date:** 2020-2-18

# **Project Overview**



**Project Name** 

Dusp5

**Project type** 

Cas9-CKO


Strain background

C57BL/6JGpt

## Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Dusp5* gene. The schematic diagram is as follows:



### Technical routes



- ➤ The *Dusp5* gene has 1 transcript. According to the structure of *Dusp5* gene, exon2 of *Dusp5-201*(ENSMUST00000038287.6) transcript is recommended as the knockout region. The region contains 149bp coding sequence.

  Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Dusp5* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

### **Notice**



- According to the existing MGI data, Mice homozygous for a knock-out allele exhibit increased proliferation and apoptosis and altered metabolic profiles in T cells. Mice homozygous for other null alleles exhibit altered eosinophilic response to parasicite infection and increased susceptibility to induced tumors.
- $\rightarrow$  The 5-terminal regulation of Gm50394 may be affectd.
- The *Dusp5* gene is located on the Chr19. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

### Gene information (NCBI)



#### Dusp5 dual specificity phosphatase 5 [ Mus musculus (house mouse) ]

Gene ID: 240672, updated on 28-Dec-2019

Summary

#### Official Symbol Dusp5 provided by MGI

Official Full Name dual specificity phosphatase 5 provided by MGI

Primary source MGI:MGI:2685183

See related Ensembl: ENSMUSG00000034765

Gene type protein coding RefSeq status VALIDATED Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae

Murinae: Mus: Mus

Also known as Gm337

Expression Broad expression in thymus adult (RPKM 25.5), duodenum adult (RPKM 24.5) and 19 other tissues See more

Orthologs human all


#### Genomic context

Location: 19: 19 D2

See Dusp5 in Genome Data Viewer

Exon count: 4

Annotation release	Status	Assembly	Chr	Location	
108	current	GRCm38.p6 (GCF_000001635.26)	19	NC_000085.6 (5352914553541809)	
Build 37.2	previous assembly	MGSCv37 (GCF_000001635.18)	19	NC_000085.5 (5360380853615812)	




# Transcript information (Ensembl)



The gene has 1 transcript, and the transcript is shown below:


Name 🍦	Transcript ID 🍦	bp 🌲	Protein	Biotype 🍦	CCDS 🍦	UniProt 🍦		Flags	± V
Dusp5-201	ENSMUST00000038287.6	2473	<u>384aa</u>	Protein coding	CCDS38024&	Q1HL35@	TSL:5	GENCODE basic	APPRIS P1

The strategy is based on the design of *Dusp5-201* transcript, The transcription is shown below




### Genomic location distribution






### Protein domain





# Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Mice homozygous for a knock-out allele exhibit increased proliferation and apoptosis and altered metabolic profiles in T cells. Mice homozygous for other null alleles exhibit altered eosinophilic response to parasicite infection and increased susceptibility to induced tumors.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





