



# ***Chst1 Cas9-CKO Strategy***

**Designer:**

**Huimin Su**

**Reviewer:**

**Ruirui Zhang**

**Design Date:**

**2020-3-12**

# Project Overview

---

**Project Name*****Chst1***

---

---

**Project type****Cas9-CKO**

---


---

**Strain background****C57BL/6JGpt**

---

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Chst1* gene. The schematic diagram is as follows:



# Technical routes

- The *Chst1* gene has 1 transcript. According to the structure of *Chst1* gene, exon3 of *Chst1-201* (ENSMUST00000065797.6) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Chst1* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.



集萃药康  
GemPharmatech

# Notice

- According to the existing MGI data, mice homozygous for a knock-out allele exhibit increased male sibling aggression when house together, a small decrease in the peripheral and mesenteric lymph nodes and peripheral blood and a small increase in the peripheral lymph nodes and peripheral blood.
- The *Chst1* gene is located on the Chr2. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

# Gene information (NCBI)

## Chst1 carbohydrate sulfotransferase 1 [*Mus musculus* (house mouse)]

Gene ID: 76969, updated on 24-Oct-2019

### Summary

Official Symbol	Chst1 provided by MGI
Official Full Name	carbohydrate sulfotransferase 1 provided by MGI
Primary source	MGI:MGI:1924219
See related	<a href="#">Ensembl:ENSMUSG00000027221</a>
Gene type	protein coding
RefSeq status	VALIDATED
Organism	<a href="#">Mus musculus</a>
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	C6ST; Gst1; KSST; GST-1; KS6ST; AW125896; KSGAL6ST; 2610008E20Rik
Expression	Broad expression in frontal lobe adult (RPKM 51.0), cortex adult (RPKM 49.6) and 16 other tissues <a href="#">See more</a>
Orthologs	<a href="#">human</a> <a href="#">all</a>


### Genomic context

Location: 2; 2 E1

[See Chst1 in Genome Data Viewer](#)

Exon count: 5

Annotation release	Status	Assembly	Chr	Location
108	current	GRCm38.p6 ( <a href="#">GCF_000001635.26</a> )	2	NC_000068.7 (92599632..92615252)
Build 37.2	previous assembly	MGSCv37 ( <a href="#">GCF_000001635.18</a> )	2	NC_000068.6 (92439864..92455409)




# Transcript information (Ensembl)


The gene has 1 transcript, and the transcript is shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Chst1-201	ENSMUST00000065797_6	2680	411aa	Protein coding	CCDS16450	Q3UY35 Q9EQC0	TSL:1 GENCODE basic APPRIS P1

The strategy is based on the design of *Chst1-201* transcript, the transcription is shown below




# Genomic location distribution



# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, mice homozygous for a knock-out allele exhibit increased male sibling aggression when house together, a small decrease in the peripheral and mesenteric lymph nodes and peripheral blood and a small increase in the peripheral lymph nodes and peripheral blood.



If you have any questions, you are welcome to inquire.

Tel: 400-9660890



集萃药康生物科技  
GemPharmatech Co.,Ltd

