

# *Cxcl17* Cas9-CKO Strategy

**Designer:**

**JiaYu**

**Reviewer:**

**Xiaojing Li**

**Design Date:**

**2020-3-25**

# Project Overview

**Project Name**

*Cxcl17*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Cxcl17* gene. The schematic diagram is as follows:



# Technical routes

- The *Cxcl17* gene has 2 transcripts. According to the structure of *Cxcl17* gene, exon1 of *Cxcl17-202* (ENSMUST00000200880.3) transcript is recommended as the knockout region. The region contains start codon ATG. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Cxcl17* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- The flox region is in the intron of the 4732471J01Rik gene, which may affect the regulation of this gene.
- The *Cxcl17* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

## Cxcl17 chemokine (C-X-C motif) ligand 17 [ *Mus musculus* (house mouse) ]

Gene ID: 232983, updated on 13-Mar-2020

### Summary


Official Symbol	Cxcl17 provided by <a href="#">MGI</a>
Official Full Name	chemokine (C-X-C motif) ligand 17 provided by <a href="#">MGI</a>
Primary source	<a href="#">MGI:MGI:2387642</a>
See related	<a href="#">Ensembl:ENSMUSG00000060188</a>
Gene type	protein coding
RefSeq status	VALIDATED
Organism	<a href="#">Mus musculus</a>
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	Vcc1; VCC-1
Expression	Biased expression in stomach adult (RPKM 37.4), lung adult (RPKM 15.8) and 3 other tissues <a href="#">See more</a>
Orthologs	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 2 transcript, and the transcript is shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Cxcl17-202	<a href="#">ENSMUST00000200880.3</a>	760	<a href="#">119aa</a>	Protein coding	<a href="#">CCDS80680</a>	<a href="#">Q5UW37</a>	TSL:1 GENCODE basic APPRIS P2
Cxcl17-201	<a href="#">ENSMUST00000074040.3</a>	761	<a href="#">128aa</a>	Protein coding	-	<a href="#">E9Q192</a>	TSL:5 GENCODE basic APPRIS ALT2

The strategy is based on the design of *Cxcl17-202* transcript, The transcription is shown below




# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

