

Capn8 Cas9-CKO Strategy

Designer:

Daohua Xu

Reviewer:

Huimin Su

Design Date:

2019-9-16

Project Overview



Project Name

Project type Cas9-CKO


Strain background C57BL/6JGpt

Capn8

Conditional Knockout strategy



This model will use CRISPR/Cas9 technology to edit the Capn8 gene. The schematic diagram is as follows:



Technical routes



- > The Capn8 gene has 4 transcripts. According to the structure of Capn8 gene, exon2 of Capn8-202

 (ENSMUST00000168514.6) transcript is recommended as the knockout region. The region contains 70bp coding sequence.

 Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Capn8* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

Notice



- > According to the existing MGI data, Mice homozygous for a knock-out allele or allele that produces a proteolytically inactive protein exhibit increased sensitivity to ethanol-induced gastric mucosa injury.
- The *Capn8* gene is located on the Chr1. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.

Gene information (NCBI)



Capn8 calpain 8 [Mus musculus (house mouse)]

Gene ID: 170725, updated on 31-Jan-2019

Summary

☆ ?

Official Symbol Capn8 provided by MGI
Official Full Name calpain 8 provided by MGI

Primary source MGI:MGI:2181366

See related Ensembl:ENSMUSG00000038599

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as nCL-2, nCL-2'

Expression Biased expression in stomach adult (RPKM 4.6), colon adult (RPKM 1.6) and 6 other tissues See more

Orthologs <u>human all</u>


Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:


Name	Transcript ID 👙	bp 🌲	Protein	Biotype	CCDS	UniProt #	Flags		
Capn8-202	ENSMUST00000168514.6	2483	381aa	Protein coding	CCDS48477 ₽	Q91VA3₽		TSL:1 GENCODE	basic
Capn8-201	ENSMUST00000048941.13	2406	703aa	Protein coding	CCDS35814 @	A0A0R4J0K6日	TSL:1	GENCODE basic	APPRIS P1
Capn8-203	ENSMUST00000192671.1	2333	690aa	Nonsense mediated decay	-	A0A0A6YVU9₽	TSL:1		
Capn8-204	ENSMUST00000193260.1	2483	No protein	IncRNA	5-	12-		TSL:1	

The strategy is based on the design of Capn8-202 transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Mice homozygous for a knock-out allele or allele that produces a proteolytically inactive protein exhibit increased sensitivity to ethanol-induced gastric mucosa injury.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





