

# *Pla2g7* Cas9-CKO Strategy

Designer: Xueting Zhang

Reviewer: Yanhua Shen

Date: 2020-03-11

# Project Overview

**Project Name**

*Pla2g7*

**Project type**


**Cas9-CKO**

**Strain background**

**C57BL/6JGpt**

# Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Pla2g7* gene. The schematic diagram is as follows:



- The *Pla2g7* gene has 6 transcripts. According to the structure of *Pla2g7* gene, exon3-exon5 of *Pla2g7-201* (ENSMUST00000024706.11) transcript is recommended as the knockout region. The region contains 361bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Pla2g7* gene. The brief process is as follows: CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.

- According to the existing MGI data, Mice homozygous for a knock-out allele exhibit reduced early mortality in response to bacterial exposure, formula feeding and asphyxia, but survivors show a significantly higher incidence of necrotizing enterocolitis relative to wild-type controls.
- Transcript *Pla2g7-203* may not be affected.
- The floxed region is near to the N-terminal of *1700071M16Rik* gene, this strategy may influence the regulatory function of the N-terminal of *1700071M16Rik* gene.
- The *Pla2g7* gene is located on the Chr17. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



# Gene information (NCBI)

**Pla2g7 phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma) [ *Mus musculus* (house mouse) ]**

Gene ID: 27226, updated on 12-Aug-2019

Summary

**Official Symbol** Pla2g7 provided by MGI  
**Official Full Name** phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma) provided by MGI  
**Primary source** MGI:MGI:1351327  
**See related** Ensembl:ENSMUSG00000023913  
**Gene type** protein coding  
**RefSeq status** VALIDATED  
**Organism** *Mus musculus*  
**Lineage** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus  
**Also known as** PAF-AH; R75400  
**Expression** Broad expression in cerebellum adult (RPKM 43.2), frontal lobe adult (RPKM 30.8) and 16 other tissues [See more](#)  
**Orthologs** [human](#) [all](#)

Genomic context

**Location:** 17; 17 B3 See Pla2g7 in [Genome Data Viewer](#)  
**Exon count:** 14


Annotation release	Status	Assembly	Chr	Location
<a href="#">108</a>	current	GRCm38.p6 ( <a href="#">GCF_000001635.26</a> )	17	NC_000083.6 (43567994..43612202)
Build 37.2	previous assembly	MGSCv37 ( <a href="#">GCF_000001635.18</a> )	17	NC_000083.5 (43705400..43749150)

# Transcript information (Ensembl)


The gene has 6 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Pla2g7-201	<a href="#">ENSMUST00000024706.11</a>	2097	<a href="#">440aa</a>	Protein coding	<a href="#">CCDS28796</a>	<a href="#">Q60963</a>	TSL:1 GENCODE basic APPRIS P1
Pla2g7-206	<a href="#">ENSMUST00000169694.1</a>	1112	<a href="#">298aa</a>	Protein coding	-	<a href="#">E9Q330</a>	CDS 3' incomplete TSL:1
Pla2g7-204	<a href="#">ENSMUST00000167214.7</a>	740	<a href="#">211aa</a>	Protein coding	-	<a href="#">E9Q6J0</a>	CDS 3' incomplete TSL:5
Pla2g7-205	<a href="#">ENSMUST00000167418.7</a>	551	<a href="#">122aa</a>	Protein coding	-	<a href="#">E9Q4T5</a>	CDS 3' incomplete TSL:3
Pla2g7-202	<a href="#">ENSMUST00000163489.1</a>	1374	No protein	Retained intron	-	-	TSL:1
Pla2g7-203	<a href="#">ENSMUST00000165706.7</a>	727	No protein	lncRNA	-	-	TSL:2

The strategy is based on the design of *Pla2g7-201* transcript,The transcription is shown below



# Genomic location distribution






# Protein domain



# Mouse phenotype description(MGI)



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Mice homozygous for a knock-out allele exhibit reduced early mortality in response to bacterial exposure, formula feeding and asphyxia, but survivors show a significantly higher incidence of necrotizing enterocolitis relative to wild-type controls.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

