

Lor Cas9-KO Strategy

Designer: Xueting Zhang

Reviewer: Yanhua Shen

Date: 2019-10-27

Project Overview

Project Name

Lor

Project type


Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Lor* gene. The schematic diagram is as follows:



- The *Lor* gene has 1 transcript. According to the structure of *Lor* gene, exon2 of *Lor-201* (ENSMUST00000058150.7) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Lor* gene. The brief process is as follows: CRISPR/Cas9 system w

- According to the existing MGI data, Mice homozygous for disruptions in this gene are runted at birth, have a translucent skin and skin skin barrier defect. The morphological skin phenotype disappears after 4-5 days.
- The *Lor* gene is located on the Chr3. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Lor loricrin [*Mus musculus* (house mouse)]

Gene ID: 16939, updated on 24-Sep-2019

Summary

- Official Symbol** Lor provided by [MGI](#)
- Official Full Name** loricrin provided by [MGI](#)
- Primary source** [MGI:MGI:96816](#)
- See related** [Ensembl:ENSMUSG00000043165](#)
- Gene type** protein coding
- RefSeq status** VALIDATED
- Organism** [Mus musculus](#)
- Lineage** Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
- Also known as** S77319; AI036317
- Expression** Restricted expression toward stomach adult (RPKM 762.9) [See more](#)
- Orthologs** [human](#) [all](#)

Genomic context

Location: 3 F1; 3 40.14 cM

See Lor in [Genome Data Viewer](#)

Exon count: 2


Annotation release	Status	Assembly	Chr	Location
108	current	GRCm38.p6 (GCF_000001635.26)	3	NC_000069.6 (92080271..92083142, complement)
Build 37.2	previous assembly	MGSCv37 (GCF_000001635.18)	3	NC_000069.5 (91884193..91887064, complement)

Transcript information (Ensembl)


The gene has 1 transcript, and the transcript is shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Lor-201	ENSMUST00000058150.7	1779	486aa	Protein coding	CCDS17546	P18165	TSL:1 GENCODE basic APPRIS P1


The strategy is based on the design of *Lor-201* transcript, The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Mice homozygous for disruptions in this gene are runted at birth, have a translucent skin and skin skin barrier defect. The morphological skin phenotype disappears after 4-5 days.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

