

Lrig1 Cas9-KO Strategy

Designer:

Yang Zeng

Reviewer:

Ruirui Zhang

Design Date:

2019-12-23

Project Overview

Project Name

Lrig1

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Lrig1* gene. The schematic diagram is as follows:



- The *Lrig1* gene has 7 transcripts. According to the structure of *Lrig1* gene, exon4-exon12 of *Lrig1-202* (ENSMUST00000101126.2) transcript is recommended as the knockout region. The region contains 1103bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Lrig1* gene. The brief process is as follows: CRISPR/Cas9 system will

- According to the existing MGI data, Homozygous null mice developed psoriasiform epidermal hyperplasia. Homozygotes exhibit hair follicle, epidermis, vertebral, eye and hearing abnormalities, decreased body size and fat amount, and increased susceptibility to bacterial infection.
- The *Lrig1* gene is located on the Chr6. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Lrig1 leucine-rich repeats and immunoglobulin-like domains 1 [*Mus musculus* (house mouse)]

Gene ID: 16206, updated on 16-Dec-2019

Summary



Official Symbol	Lrig1 provided by MGI
Official Full Name	leucine-rich repeats and immunoglobulin-like domains 1 provided by MGI
Primary source	MGI:MGI:107935
See related	Ensembl:ENSMUSG00000030029
Gene type	protein coding
RefSeq status	VALIDATED
Organism	Mus musculus
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	Img; LIG-1; D6Bwg0781e
Expression	Ubiquitous expression in colon adult (RPKM 30.2), cerebellum adult (RPKM 20.8) and 26 other tissues See more
Orthologs	human all

Transcript information (Ensembl)


The gene has 7 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Translation ID	Biotype	CCDS	UniProt	Flags
Lrig1-202	ENSMUST000000101126.2	4951	1091aa	ENSMUSP00000098686.2	Protein coding	CCDS20380	P70193	TSL:1 GENCODE basic APPRIS P1
Lrig1-201	ENSMUST00000032105.10	4831	1091aa	ENSMUSP00000032105.4	Protein coding	CCDS20380	P70193	TSL:1 GENCODE basic APPRIS P1
Lrig1-207	ENSMUST000000204645.2	4849	1091aa	ENSMUSP000000144963.1	Nonsense mediated decay	-	P70193	TSL:1
Lrig1-205	ENSMUST000000150811.7	700	No protein	-	lncRNA	-	-	TSL:3
Lrig1-204	ENSMUST000000141014.7	666	No protein	-	lncRNA	-	-	TSL:3
Lrig1-203	ENSMUST000000124165.1	587	No protein	-	lncRNA	-	-	TSL:3
Lrig1-206	ENSMUST000000203876.1	438	No protein	-	lncRNA	-	-	TSL:3


The strategy is based on the design of *Lrig1*-202 transcript,The transcription is shown below



Genomic location distribution




Protein domain



Mouse phenotype description(MGI)

Phenotype Overview ?



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Homozygous null mice developed psoriasiform epidermal hyperplasia. Homozygotes exhibit hair follicle, epidermis, vertebral, eye and hearing abnormalities, decreased body size and fat amount, and increased susceptibility to bacterial infection.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

