

Trpv5 Cas9-KO Strategy

Designer: Yang Zeng **Reviewer:**

Xiaojing Li

Design Date:

2019-11-26

Project Overview



Project Name

Trpv5

Project type

Cas9-KO


Strain background

C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Trpv5* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Trpv5* gene has 2 transcripts. According to the structure of *Trpv5* gene, exon2-exon7 of *Trpv5-201* (ENSMUST00000031901.10) transcript is recommended as the knockout region. The region contains 778bp coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Trpv5* gene. The brief process is as follows: CRISPR/Cas9 system we

Notice



- ➤ According to the existing MGI data, Homozygous mutant mice exhibit increased calcium excretion and reduced bone thickenss.
- > The *Trpv5* gene is located on the Chr6. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Trpv5 transient receptor potential cation channel, subfamily V, member 5 [Mus musculus (house mouse)]

Gene ID: 194352, updated on 25-Nov-2019

Summary



Official Symbol Trpv5 provided by MGI

Official Full Name transient receptor potential cation channel, subfamily V, member 5 provided by MGI

Primary source MGI:MGI:2429764

> Ensembl: ENSMUSG00000036899 See related

Gene type protein coding RefSeq status PROVISIONAL

Organism Mus musculus


Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as CAT2; ECAC1; D630033B11

Expression Biased expression in kidney adult (RPKM 1.9), placenta adult (RPKM 0.2) and 2 other tissues See more

Orthologs <u>human</u> all




Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:


Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Trpv5-201	ENSMUST00000031901.10	3243	723aa	Protein coding	CCDS20053	P69744	TSL:1 GENCODE basic APPRIS P2
Trpv5-202	ENSMUST00000193503.1	2025	674aa	Protein coding	8+	A0A0A6YW69	TSL:5 GENCODE basic APPRIS ALT2

The strategy is based on the design of *Trpv5-201* transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, Homozygous mutant mice exhibit increased calcium excretion and reduced bone thickenss.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





