

Deptor Cas9-KO Strategy

Designer: Xiaojing Li

Design Date: 2020-1-22

Reviewer: JiaYu

Project Overview



Project Name Deptor


Project type Cas9-KO

Strain background C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Deptor* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Deptor* gene has 4 transcripts. According to the structure of *Deptor* gene, exon2 of *Deptor-202*(ENSMUST00000096433.9) transcript is recommended as the knockout region. The region contains 179bp coding sequence.

 Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Deptor* gene. The brief process is as follows: CRISPR/Cas9 system

Notice



- > The *Deptor* gene is located on the Chr15. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- > This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Deptor DEP domain containing MTOR-interacting protein [Mus musculus (house mouse)]

Gene ID: 97998, updated on 8-Dec-2019

Summary

Official Symbol Deptor provided by MGI

Official Full Name DEP domain containing MTOR-interacting protein provided by MGI

Primary source MGI:MGI:2146322

See related Ensembl: ENSMUSG00000022419

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as Depdc6

Expression Broad expression in bladder adult (RPKM 11.6), subcutaneous fat pad adult (RPKM 8.1) and 23 other tissues See more

Orthologs human all


Transcript information (Ensembl)



The gene has 4 transcripts, all transcripts are shown below:


| and the second | offine. | | | | | | |
|----------------|-----------------------|------|--------------|-------------------------|-----------|---------------|-------------------------------|
| Name | Transcript ID | bp | Protein | Biotype | CCDS | UniProt | Flags |
| Deptor-202 | ENSMUST00000096433.9 | 8547 | 409aa | Protein coding | CCDS27474 | B2ZRS3 Q570Y9 | TSL:1 GENCODE basic APPRIS P1 |
| Deptor-201 | ENSMUST00000023056.7 | 3300 | <u>397aa</u> | Protein coding | CCDS27475 | B2ZRS7 Q570Y9 | TSL:1 GENCODE basic |
| Deptor-203 | ENSMUST00000100660.10 | 1377 | <u>401aa</u> | Protein coding | - | B2ZRS5 | TSL:1 GENCODE basic |
| Deptor-204 | ENSMUST00000226687.1 | 693 | 100aa | Nonsense mediated decay | - | A0A2I3BRE1 | |

The strategy is based on the design of Deptor-202 transcript, The transcription is shown below




Genomic location distribution





Protein domain







If you have any questions, you are welcome to inquire. Tel: 400-9660890





