



Pla2g3 Cas9-CKO Strategy

Designer: Jia Yu

Reviewer: Xiaojing Li

Design Date: 2020-8-13

Project Overview


Project Name***Pla2g3***

Project type**Cas9-CKO**

Strain background**C57BL/6JGpt**

Conditional Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Pla2g3* gene. The schematic diagram is as follows:



Technical routes

- The *Pla2g3* gene has 3 transcripts. According to the structure of *Pla2g3* gene, exon2-exon6 of *Pla2g3*-202(ENSMUST00000064265.12) transcript is recommended as the knockout region. The region contains 799bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Pla2g3* gene. The brief process is as follows:CRISPR/Cas9 system and Donor were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.
- The flox mice will be knocked out after mating with mice expressing Cre recombinase, resulting in the loss of function of the target gene in specific tissues and cell types.



集萃药康
GemPharmatech

Notice

- According to the existing MGI data, homozygous mice have reduced male fertility with smaller litter size and decreased susceptibility to IgE antigen-induced passive systemic and cutaneous anaphylaxis due to impaired mast cell maturation, degranulation and histamine release.
- The *Pla2g3* gene is located on the Chr11. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of loxp insertion on gene transcription, RNA splicing and protein translation cannot be predicted at existing technological level.



集萃药康
GemPharmatech

Gene information (NCBI)

Pla2g3 phospholipase A2, group III [Mus musculus (house mouse)]

Gene ID: 237625, updated on 13-Mar-2020

Summary



Official Symbol Pla2g3 provided by [MGI](#)

Official Full Name phospholipase A2, group III provided by [MGI](#)

Primary source [MGI:MG1:2444945](#)

See related [Ensembl:ENSMUSG00000034579](#)

Gene type protein coding

RefSeq status VALIDATED

Organism [Mus musculus](#)

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Expression Broad expression in colon adult (RPKM 2.6), frontal lobe adult (RPKM 2.2) and 18 other tissues [See more](#)


Orthologs [human](#) [all](#)

Transcript information (Ensembl)


The gene has 3 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Pla2g3-202	ENSMUST00000064265.12	2566	504aa	Protein coding	CCDS48740	Q8BZT7	TSL:1 GENCODE basic APPRIS P1
Pla2g3-201	ENSMUST00000044682.6	2608	446aa	Nonsense mediated decay	-	Q8BV23	TSL:2
Pla2g3-203	ENSMUST00000151860.1	2869	No protein	Retained intron	-	-	TSL:5

The strategy is based on the design of *Pla2g3-202* transcript, the transcription is shown below:



Genomic location distribution



Protein domain

ENSMUSP000000068...

MobiDB lite

Low complexity (Seq)

Cleavage site (Sign...)

Superfamily

Pfam

PROSITE patterns

PANTHER

PTHR12253

PTHR12253:SF19

Gene3D

CDD

All sequence SNPs/i...

Sequence variants (dbSNP and all other sources)


Variant Legend

- inframe deletion
- missense variant
- splice region variant
- synonymous variant

Scale bar

0 60 120 180 240 300 360 420 480 504

Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, homozygous mice have reduced male fertility with smaller litter size and decreased susceptibility to IgE antigen-induced passive systemic and cutaneous anaphylaxis due to impaired mast cell maturation, degranulation and histamine release.



If you have any questions, you are welcome to inquire.

Tel: 400-9660890



集萃药康生物科技
GemPharmatech Co.,Ltd

