

# *Pde8a* Cas9-KO Strategy

**Designer:**

**JiaYu**

**Reviewer:**

**Xiaojing Li**

**Design Date:**

**2020-2-11**

# Project Overview

**Project Name**

*Pde8a*

**Project type**


**Cas9-KO**

**Strain background**

**C57BL/6JGpt**

# Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Pde8a* gene. The schematic diagram is as follows:



- The *Pde8a* gene has 3 transcripts. According to the structure of *Pde8a* gene, exon2-exon3 of *Pde8a-201* (ENSMUST00000026672.7) transcript is recommended as the knockout region. The region contains 248bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Pde8a* gene. The brief process is as follows: CRISPR/Cas9 system

- According to the existing MGI data, Targeted disruption of this gene results in a 4-fold increase in basal release of testosterone in isolated Leydig cells as well as a significant increase in the sensitivity to luteinizing hormone, measured as testosterone released into the media.
- The *Pde8a* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

# Gene information (NCBI)

## Pde8a phosphodiesterase 8A [Mus musculus (house mouse)]

Gene ID: 18584, updated on 31-Jan-2019

### Summary




<b>Official Symbol</b>	Pde8a provided by <a href="#">MGI</a>
<b>Official Full Name</b>	phosphodiesterase 8A provided by <a href="#">MGI</a>
<b>Primary source</b>	<a href="#">MGI:MGI:1277116</a>
<b>See related</b>	<a href="#">Ensembl:ENSMUSG00000025584</a>
<b>Gene type</b>	protein coding
<b>RefSeq status</b>	VALIDATED
<b>Organism</b>	<a href="#">Mus musculus</a>
<b>Lineage</b>	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
<b>Also known as</b>	AI551852, Pde8
<b>Expression</b>	Broad expression in testis adult (RPKM 48.2), kidney adult (RPKM 13.7) and 18 other tissues <a href="#">See more</a>
<b>Orthologs</b>	<a href="#">human</a> <a href="#">all</a>

# Transcript information (Ensembl)


The gene has 3 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Pde8a-201	<a href="#">ENSMUST00000026672.7</a>	3802	<a href="#">823aa</a>	Protein coding	<a href="#">CCDS40005</a>	<a href="#">Q88502</a>	TSL:1 GENCODE basic APPRIS P1
Pde8a-203	<a href="#">ENSMUST00000130494.1</a>	861	No protein	lncRNA	-	-	TSL:3
Pde8a-202	<a href="#">ENSMUST00000128154.7</a>	583	No protein	lncRNA	-	-	TSL:3


The strategy is based on the design of *Pde8a-201* transcript,The transcription is shown below




# Genomic location distribution



# Protein domain



# Mouse phenotype description(MGI )



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Targeted disruption of this gene results in a 4-fold increase in basal release of testosterone in isolated Leydig cells as well as a significant increase in the sensitivity to luteinizing hormone, measured as testosterone released into the media.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

