

# *Slc22a18* Cas9-KO Strategy

**Designer:**

**JiaYu**

**Reviewer:**

**Xiaojing Li**

**Design Date:**

**2020-2-11**

# Project Overview

**Project Name**

*Slc22a18*

**Project type**

**Cas9-KO**

**Strain background**

**C57BL/6JGpt**

# Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Slc22a18* gene. The schematic diagram is as follows:



- The *Slc22a18* gene has 5 transcripts. According to the structure of *Slc22a18* gene, exon3-exon5 of *Slc22a18-202* (ENSMUST00000105917.2) transcript is recommended as the knockout region. The region contains 392bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Slc22a18* gene. The brief process is as follows: CRISPR/Cas9 syst

- The *Slc22a18* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

# Gene information (NCBI)

## Slc22a18 solute carrier family 22 (organic cation transporter), member 18 [Mus musculus (house mouse)]

Gene ID: 18400, updated on 19-Mar-2019

### Summary



|                           |                                                                                                                                                                           |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Official Symbol</b>    | Slc22a18 provided by <a href="#">MGI</a>                                                                                                                                  |
| <b>Official Full Name</b> | solute carrier family 22 (organic cation transporter), member 18 provided by <a href="#">MGI</a>                                                                          |
| <b>Primary source</b>     | <a href="#">MGI:MGI:1336884</a>                                                                                                                                           |
| <b>See related</b>        | <a href="#">Ensembl:ENSMUSG000000000154</a>                                                                                                                               |
| <b>Gene type</b>          | protein coding                                                                                                                                                            |
| <b>RefSeq status</b>      | VALIDATED                                                                                                                                                                 |
| <b>Organism</b>           | <a href="#">Mus musculus</a>                                                                                                                                              |
| <b>Lineage</b>            | Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus |
| <b>Also known as</b>      | AW260131, BWR1A, BWSCR1A, HET, ITM, Impt1, Orct12, Slc22a1l, TSSC5, p45-BWR1A                                                                                             |
| <b>Expression</b>         | Biased expression in kidney adult (RPKM 388.3), duodenum adult (RPKM 122.8) and 5 other tissues <a href="#">See more</a>                                                  |
| <b>Orthologs</b>          | <a href="#">human</a> <a href="#">all</a>                                                                                                                                 |

# Transcript information (Ensembl)


The gene has 5 transcripts,all transcripts are shown below:

| Name         | Transcript ID                         | bp   | Protein               | Biotype                 | CCDS                      | UniProt                    | Flags                         |
|--------------|---------------------------------------|------|-----------------------|-------------------------|---------------------------|----------------------------|-------------------------------|
| Slc22a18-202 | <a href="#">ENSMUST00000105917.2</a>  | 1552 | <a href="#">410aa</a> | Protein coding          | <a href="#">CCDS22041</a> | <a href="#">A0A0R4J0P7</a> | TSL:1 GENCODE basic APPRIS P1 |
| Slc22a18-201 | <a href="#">ENSMUST00000052348.11</a> | 1525 | <a href="#">410aa</a> | Protein coding          | <a href="#">CCDS22041</a> | <a href="#">A0A0R4J0P7</a> | TSL:1 GENCODE basic APPRIS P1 |
| Slc22a18-204 | <a href="#">ENSMUST00000145943.7</a>  | 1383 | <a href="#">65aa</a>  | Nonsense mediated decay | -                         | <a href="#">D6RGN1</a>     | TSL:1                         |
| Slc22a18-203 | <a href="#">ENSMUST00000141988.7</a>  | 862  | <a href="#">44aa</a>  | Nonsense mediated decay | -                         | <a href="#">D6RDC9</a>     | TSL:3                         |
| Slc22a18-205 | <a href="#">ENSMUST00000150791.7</a>  | 862  | <a href="#">44aa</a>  | Nonsense mediated decay | -                         | <a href="#">D6RDC9</a>     | TSL:5                         |


The strategy is based on the design of *Slc22a18-202* transcript,The transcription is shown below



# Genomic location distribution



# Protein domain



If you have any questions, you are welcome to inquire.

Tel: 400-9660890

