

Aacs Cas9-KO Strategy

Designer: Ruirui Zhang

Reviewer: Huimin Su

Design Date: 2020-2-13

Project Overview



Project Name

Aacs

Project type

Cas9-KO


Strain background

C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the Aacs gene. The schematic diagram is as follows:



Technical routes



- ➤ The Aacs gene has 6 transcripts. According to the structure of Aacs gene, exon2 of Aacs-201

 (ENSMUST00000031445.4) transcript is recommended as the knockout region. The region contains 104bp coding sequence.

 Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Aacs* gene. The brief process is as follows: CRISPR/Cas9 system v

Notice



- > The *Aacs* gene is located on the Chr5. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Aacs acetoacetyl-CoA synthetase [Mus musculus (house mouse)]

Gene ID: 78894, updated on 12-Aug-2019

Summary

☆ ?

Official Symbol Aacs provided by MGI

Official Full Name acetoacetyl-CoA synthetase provided by MGI

Primary source MGI:MGI:1926144

See related Ensembl: ENSMUSG00000029482

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae;

Murinae; Mus; Mus

Also known as SUR5; 2210408B16Rik

Expression Broad expression in mammary gland adult (RPKM 107.7), subcutaneous fat pad adult (RPKM 90.4) and 27 other tissues See more

Orthologs human all


Transcript information (Ensembl)



The gene has 6 transcripts, all transcripts are shown below:


Name 4	Transcript ID	bp 🌲	Protein	Biotype A	CCDS 🍦	UniProt 4	Flags
Aacs-203	ENSMUST00000144361.1	796	No protein	Retained intron	171	67.4	TSL:5
Aacs-202	ENSMUST00000131355.1	527	No protein	Retained intron	120	(4.5)	TSL:2
Aacs-201	ENSMUST00000031445.4	3239	<u>672aa</u>	Protein coding	CCDS19687 ₽	Q9D2R0 ₺	TSL:1 GENCODE basic APPRIS P1
Aacs-206	ENSMUST00000200286.1	245	No protein	Processed transcript	24		TSL:5
Aacs-204	ENSMUST00000196556.1	163	No protein	Processed transcript	876	(7.0	TSL:1
Aacs-205	ENSMUST00000199978.1	158	No protein	Processed transcript	©±0	823	TSL:1

The strategy is based on the design of Aacs-201 transcript, the transcription is shown below:




Genomic location distribution





Protein domain







If you have any questions, you are welcome to inquire. Tel: 400-9660890





