

Plaa Cas9-KO Strategy

Designer:

JiaYu

Reviewer:

Xiaojing Li

Design Date:

2020-2-18

Project Overview

Project Name

Plaa

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Plaa* gene. The schematic diagram is as follows:



- The *Plaa* gene has 6 transcripts. According to the structure of *Plaa* gene, exon2-exon9 of *Plaa-201* (ENSMUST00000107107.8) transcript is recommended as the knockout region. The region contains 1268bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Plaa* gene. The brief process is as follows: CRISPR/Cas9 system v

- According to the existing MGI data, Homozygous KO is embryonic lethal. A hypomorphic homozygous point mutation affects neuromuscular junctions and Purkinje cell development, causing early-onset neurodysfunction.
- The *Plaa* gene is located on the Chr4. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)

Plaa phospholipase A2, activating protein [Mus musculus (house mouse)]

Gene ID: 18786, updated on 31-Jan-2019

Summary




Official Symbol	Plaa provided by MGI
Official Full Name	phospholipase A2, activating protein provided by MGI
Primary source	MGI:MGI:104810
See related	Ensembl:ENSMUSG00000028577
Gene type	protein coding
RefSeq status	VALIDATED
Organism	Mus musculus
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus
Also known as	2410007N06, AI536418, AU018445, AW208417, D4Ert618e, PLA2P, PLAP, Ufd3
Expression	Ubiquitous expression in CNS E11.5 (RPKM 9.7), placenta adult (RPKM 8.4) and 28 other tissues See more
Orthologs	human all

Transcript information (Ensembl)


The gene has 6 transcripts,all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Plaa-201	ENSMUST00000107107.8	2784	794aa	Protein coding	CCDS18359	P27612	TSL:1 GENCODE basic APPRIS P1
Plaa-202	ENSMUST00000127656.2	641	214aa	Protein coding	-	F7D1R5	5' and 3' truncations in transcript evidence prevent annotation of the start and the end of the CDS. CDS 5' and 3' incomplete TSL:2
Plaa-203	ENSMUST00000129748.1	3886	No protein	lncRNA	-	-	TSL:2
Plaa-205	ENSMUST00000136669.1	1514	No protein	lncRNA	-	-	TSL:1
Plaa-206	ENSMUST00000146118.1	693	No protein	lncRNA	-	-	TSL:2
Plaa-204	ENSMUST00000135696.1	645	No protein	lncRNA	-	-	TSL:3


The strategy is based on the design of *Plaa-201* transcript,The transcription is shown below




Genomic location distribution



Protein domain



Mouse phenotype description(MGI)



Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).

According to the existing MGI data, Homozygous KO is embryonic lethal. A hypomorphic homozygous point mutation affects neuromuscular junctions and Purkinje cell development, causing early-onset neurodysfunction.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

