

Nprl3 Cas9-KO Strategy

Designer:

Daohua Xu

Reviewer:

Huimin Su

Design Date:

2020-2-25

Project Overview



Project Name

Nprl3

Project type

Cas9-KO


Strain background

C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Nprl3* gene. The schematic diagram is as follows:



Technical routes



- ➤ The *Nprl3* gene has 16 transcripts. According to the structure of *Nprl3* gene, exon2-exon4 of *Nprl3-201*(ENSMUST00000020530.11) transcript is recommended as the knockout region. The region contains 275bp coding sequence Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Nprl3* gene. The brief process is as follows: CRISPR/Cas9 system

Notice



- > According to the existing MGI data, This gene is deleted in the Hbath-J mutation.
- The *Nprl3* gene is located on the Chr11. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Nprl3 nitrogen permease regulator-like 3 [Mus musculus (house mouse)]

Gene ID: 17168, updated on 31-Jan-2019

Summary

☆ ?

Official Symbol Nprl3 provided by MGI

Official Full Name nitrogen permease regulator-like 3 provided by MGI

Primary source MGI:MGI:109258

See related Ensembl:ENSMUSG00000020289

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha;

Muroidea; Muridae; Murinae; Mus; Mus

Also known as Aag, CGTHBA, HS-26, HS-40, Mare, Phg, Prox1, m(alpha)RE

Expression Ubiquitous expression in adrenal adult (RPKM 39.2), ovary adult (RPKM 30.3) and 28 other tissues See more

Orthologs <u>human</u> all


Transcript information (Ensembl)



The gene has 16 transcripts, all transcripts are shown below:


Biotype CCDS UniProt Flags tein coding CCDS24521 Q8VIJ8 TSL:1 GENCODE basic APPRIS P tein coding - A7M7S2 TSL:5 GENCODE basic tein coding - A2AAX8 CDS 3' incomplete TSL:3 e mediated decay - F2Z3Y4 TSL:1 e mediated decay - F2Z404 TSL:5
tein coding - A7M7S2 TSL:5 GENCODE basic tein coding - A2AAX8 CDS 3' incomplete TSL:3 e mediated decay - F2Z3Y4 TSL:1 e mediated decay - F2Z404 TSL:5
tein coding - A2AAX8 CDS 3' incomplete TSL:3 e mediated decay - F2Z3Y4 TSL:1 e mediated decay - F2Z404 TSL:5
e mediated decay - F2Z3Y4 TSL:1 e mediated decay - F2Z404 TSL:5
e mediated decay - F2Z404 TSL:5
The second secon
mediated decay - <u>F2Z3V7</u> TSL:1
mediated decay - <u>F2Z3V7</u> TSL:5
mediated decay - <u>D6RGB2</u> TSL:5
ained intron TSL:3
ained intron - TSL:3
ained intron - TSL:3
ained intron - TSL:2
IncRNA TSL:1
IncRNA TSL:3
IncRNA TSL:3
IncRNA TSL:2

The strategy is based on the design of Nprl3-201 transcript, The transcription is shown below




Genomic location distribution






Protein domain





Mouse phenotype description(MGI)





Phenotypes affected by the gene are marked in blue.Data quoted from MGI database(http://www.informatics.jax.org/).

According to the existing MGI data, This gene is deleted in the Hbath-J mutation.



If you have any questions, you are welcome to inquire. Tel: 400-9660890





