

# *Nt5c1a* Cas9-KO Strategy

**Designer:** Daohua Xu

**Reviewer:** Huimin Su

**Design Date:** 2020-5-20

# Project Overview

**Project Name**

*Nt5c1a*

**Project type**


**Cas9-KO**

**Strain background**

**C57BL/6JGpt**

# Knockout strategy

This model will use CRISPR/Cas9 technology to edit the *Nt5c1a* gene. The schematic diagram is as follows:



- The *Nt5c1a* gene has 1 transcript. According to the structure of *Nt5c1a* gene, exon2-exon4 of *Nt5c1a-201* (ENSMUST00000068262.5) transcript is recommended as the knockout region. The region contains 421bp coding sequence. Knock out the region will result in disruption of protein function.
- In this project we use CRISPR/Cas9 technology to modify *Nt5c1a* gene. The brief process is as follows: CRISPR/Cas9 system

- According to the existing MGI data, mice homozygous for a null allele exhibit reduced adenosine levels in muscles post-electrical stimulation.
- The *Nt5c1a* gene is located on the Chr4. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This Strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

# Gene information (NCBI)

## Nt5c1a 5'-nucleotidase, cytosolic 1A [Mus musculus (house mouse)]

Gene ID: 230718, updated on 13-Mar-2020

### Summary




|                           |                                                                                                                                                                           |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Official Symbol</b>    | Nt5c1a provided by <a href="#">MGI</a>                                                                                                                                    |
| <b>Official Full Name</b> | 5'-nucleotidase, cytosolic 1A provided by <a href="#">MGI</a>                                                                                                             |
| <b>Primary source</b>     | <a href="#">MGI:MGI:2155700</a>                                                                                                                                           |
| <b>See related</b>        | <a href="#">Ensembl:ENSMUSG00000054958</a>                                                                                                                                |
| <b>Gene type</b>          | protein coding                                                                                                                                                            |
| <b>RefSeq status</b>      | VALIDATED                                                                                                                                                                 |
| <b>Organism</b>           | <a href="#">Mus musculus</a>                                                                                                                                              |
| <b>Lineage</b>            | Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia; Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus |
| <b>Also known as</b>      | Cn1a, Gm49                                                                                                                                                                |
| <b>Expression</b>         | Biased expression in heart adult (RPKM 8.9), adrenal adult (RPKM 7.9) and 5 other tissues <a href="#">See more</a>                                                        |
| <b>Orthologs</b>          | <a href="#">human</a> <a href="#">all</a>                                                                                                                                 |

# Transcript information (Ensembl)


The gene has 1 transcript, and the transcript is shown below:

| Name       | Transcript ID                        | bp   | Protein               | Biotype        | CCDS                       | UniProt                | Flags                                                                                                                                                                                                         |
|------------|--------------------------------------|------|-----------------------|----------------|----------------------------|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Nt5c1a-201 | <a href="#">ENSMUST00000068262.5</a> | 1220 | <a href="#">365aa</a> | Protein coding | <a href="#">CCDS3887.1</a> | <a href="#">A3KFX0</a> | TSL:1 GENCODE basic APPRIS is a system to annotate alternatively spliced transcripts based on a range of computational methods to identify the most functionally important transcript(s) of a gene. APPRIS P1 |


The strategy is based on the design of *Nt5c1a-201* transcript, The transcription is shown below




# Genomic location distribution



# Protein domain



# Mouse phenotype description(MGI )



*Phenotypes affected by the gene are marked in blue. Data quoted from MGI database(<http://www.informatics.jax.org/>).*

According to the existing MGI data, Mice homozygous for a null allele exhibit reduced adenosine levels in muscles post-electrical stimulation.

If you have any questions, you are welcome to inquire.

Tel: 400-9660890

