

Olfr66 Cas9-KO Strategy

Designer: Xueting Zhang

Reviewer: Daohua Xu

Design Date: 2020-9-14

Project Overview



Project Name

Olfr66

Project type

Cas9-KO

Strain background

C57BL/6JGpt

Knockout strategy



This model will use CRISPR/Cas9 technology to edit the *Olfr66* gene. The schematic diagram is as follows:



Technical routes



- > The *Olfr66* gene has 2 transcripts. According to the structure of *Olfr66* gene, exon1 of *Olfr66*201(ENSMUST00000079117.1) transcript is recommended as the knockout region. The region contains all of the coding sequence. Knock out the region will result in disruption of protein function.
- ➤ In this project we use CRISPR/Cas9 technology to modify *Olfr66* gene. The brief process is as follows: CRISPR/Cas9 system were microinjected into the fertilized eggs of C57BL/6JGpt mice. Fertilized eggs were transplanted to obtain positive F0 mice which were confirmed by PCR and sequencing. A stable F1 generation mouse model was obtained by mating positive F0 generation mice with C57BL/6JGpt mice.

Notice



- > Dnajc19-ps gene will be deleted together in this strategy.
- > The *Olfr66* gene is located on the Chr7. If the knockout mice are crossed with other mice strains to obtain double gene positive homozygous mouse offspring, please avoid the two genes on the same chromosome.
- This strategy is designed based on genetic information in existing databases. Due to the complexity of biological processes, all risk of the gene knockout on gene transcription, RNA splicing and protein translation cannot be predicted at the existing technology level.

Gene information (NCBI)



Olfr66 olfactory receptor 66 [Mus musculus (house mouse)]

Gene ID: 18367, updated on 13-Mar-2020

Summary

☆ ?

Official Symbol Olfr66 provided by MGI

Official Full Name olfactory receptor 66 provided by MGI

Primary source MGI:MGI:1341906

See related Ensembl:ENSMUSG00000058200

Gene type protein coding
RefSeq status VALIDATED
Organism Mus musculus

Lineage Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Glires; Rodentia;

Myomorpha; Muroidea; Muridae; Murinae; Mus; Mus

Also known as 5'[b]1, 5'beta1, MOR1-3, ORL532

Summary Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of

a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and

proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008]

Orthologs human all

Transcript information (Ensembl)



The gene has 2 transcripts, all transcripts are shown below:

Name	Transcript ID	bp	Protein	Biotype	CCDS	UniProt	Flags
Olfr66-202	ENSMUST00000216303.1	3080	<u>311aa</u>	Protein coding	CCDS21594	F8VQ01	TSL:5 GENCODE basic APPRIS P1
Olfr66-201	ENSMUST00000079117.1	936	<u>311aa</u>	Protein coding	CCDS21594	F8VQ01	TSL:NA GENCODE basic APPRIS P1


The strategy is based on the design of *Olfr66-201* transcript, the transcription is shown below:

< Olfr66-201
protein coding

Reverse strand — 936 bp —


Genomic location distribution





Protein domain







If you have any questions, you are welcome to inquire. Tel: 400-9660890





